RUSH: Theriogenology article for proofing (# 11984)

Dear Author:

The proof of your article to be published by Elsevier in Theriogenology, is available as a PDF file at the following URL:

<http://rapidproof.cadmus.com/RapidProof/retrieval/index.jsp>

Login: your e-mail address

Password: HERk4k377Dw6

Also attached are instructions on the annotation of PDF files, as well as a Query Form if we have any questions regarding your article.

Proof corrections can now be annotated on-screen, which allows you to indicate your correction directly on the PDF proof. Annotating and returning your PDF proof as an email attachment will ensure quick and accurate publication of your article. To read these files and annotate the electronic proof with corrections you will need the latest version of Adobe Reader, which is available for free at:

<http://www.adobe.com/products/acrobat/readstep.html>. Please also see the accompanying instructions for further information.

If PDF annotations are not feasible, please consider one of the following methods of returning proof corrections:

- * List the corrections (including replies to any author queries) in an e-mail and return to Lorraine Bernazzani at Lbernazz@optonline.net or simply use the Reply button. Using this option, please refer to the line numbers on the proof.
- * Mark corrections and any other comments (including replies to any author queries) on a printout of the PDF file and fax it to Lorraine Bernazzani (fax # 845-883-5682).
- * If none of the above options are possible, please express mail a marked up copy of your article to the address below.

After accessing the PDF proof, please:

- * carefully proofread the entire article, including tables, equations, figure legends, and references;
- * confirm that all references, tables, and figures match up correctly with their citations in text;
- * ensure accuracy and spelling of affiliations, addresses, and author names;
- * check that any Greek letters (such as "mu") have been translated correctly;
- * verify scientific notations, drug dosages, and manufacturer names and locations;
- * be sure permission has been procured for any reprinted materials; and
- * answer all author queries completely (these are listed on the last page).

Changes that significantly alter the content of the article, such as new figures and tables or rewritten sections, will only be considered at this stage with the Editor's approval. It is important that all of your corrections are sent back to us in one communication. Please check your proof carefully before replying, as inclusion of any subsequent corrections cannot be guaranteed and there will be no further opportunity to proofread your article.

If you submitted usable color figures with your article they will appear in color on the web, at no extra

charge. In the printed issue, color reproduction depends on journal policy and whether you agree to bear any costs. Any 'supplementary' material to your article (i.e., not appearing in print) will be accessible after your corrected article is placed online; such material is not part of the proofing process and is therefore not attached here.

PLEASE RESPOND WITHIN 48 HOURS, even if you have no corrections, and include the journal name and article number on any correspondence.

Kind regards,

Lorraine Bernazzani,
Journal Manager
Elsevier Inc.
236 Patura Road
Modena, NY 12548
Phone: 845-883-5681
Fax: 845-883-5682
E-mail: Lbernazz@optonline.net

INSTRUCTIONS ON THE ANNOTATION OF PDF FILES

To view, print and annotate your article you will need Adobe Reader version 7 (or higher). This program is freely available for a whole series of platforms that include PC, Mac, and UNIX and can be downloaded from <http://get.adobe.com/reader/>. The exact system requirements are given at the Adobe site: <http://www.adobe.com/products/reader/tech-specs.html>.

PDF ANNOTATIONS	
Adobe Reader version 7, 8 or 9	Adobe Reader version X
<p>To make annotations in the PDF file, go to the main Adobe tool bar and change the cursor from a hand symbol to the normal cursor by clicking on the 'Select' button in the menu bar at the top (versions 7 and 8).</p> <p>When you open the PDF file using Adobe Reader, the Commenting tool bar should be displayed automatically; if not, click on 'Tools', select 'Comment & Markup' (or 'Commenting' in version 7), then click on 'Show Comment & Markup tool bar' (or 'Commenting tool bar' in version 7, or 'Show Commenting bar' on the Mac). If these options are not available in your Adobe Reader menus then it is possible that your Adobe Acrobat version is lower than 7 or the PDF has not been prepared properly.</p> <p>(PC, Adobe Reader version 7)</p> <p>(PC, Adobe Reader version 8, right-click on title bar (Comment & Markup) to show additional icons)</p> <p>(Mac)</p> <p>PDF ANNOTATIONS (Adobe Reader version 9)</p> <p>The default for the Commenting tool bar is set to 'off' in version 9. To change this setting select 'Edit Preferences', then 'Documents' (at left under 'Categories'), then select the option 'Never' for 'PDF/A View Mode' (the Commenting tool bar is the same as in version 8).</p> <p>(Changing the default setting, Adobe version 9)</p>	<p>To make annotations in the PDF file, open the PDF file using Adobe Reader X, click on 'Comment'.</p> <p>If this option is not available in your Adobe Reader menus then it is possible that your Adobe Acrobat version is lower than X or the PDF has not been prepared properly.</p> <p>This opens a task pane and, below that, a list of all Comments in the text. These comments initially show all the changes made by our copyeditor to your file.</p>

HOW TO...		
Action	Adobe Reader version 7, 8 or 9	Adobe Reader version X
Insert text	<p>Click the 'Text Edits' button on the Commenting tool bar. Click to set the cursor location in the text and simply start typing. The text will appear in a commenting box. You may also cut-and-paste text from another file into the commenting box. Close the box by clicking on 'x' in the top right-hand corner.</p>	<p>Click the 'Insert Text' icon on the Comment tool bar. Click to set the cursor location in the text and simply start typing. The text will appear in a commenting box. You may also cut-and-paste text from another file into the commenting box. Close the box by clicking on 'x' in the top right-hand corner.</p>
Replace text	<p>Click the 'Text Edits' button on the Commenting tool bar. To highlight the text to be replaced, click and drag the cursor over the text. Then simply type in the replacement text. The replacement text will appear in a commenting box. You may also cut-and-paste text from another file into this box. To replace formatted text (an equation for example) please Attach a file (see below).</p>	<p>Click the 'Replace (Ins)' icon on the Comment tool bar. To highlight the text to be replaced, click and drag the cursor over the text. Then simply type in the replacement text. The replacement text will appear in a commenting box. You may also cut-and-paste text from another file into this box. To replace formatted text (an equation for example) please Attach a file (see below).</p>
Remove text	<p>Click the 'Text Edits' button on the Commenting tool bar. Click and drag over the text to be deleted. Then press the delete button on your keyboard. The text to be deleted will then be struck through.</p>	<p>Click the 'Strikethrough (Del)' icon on the Comment tool bar. Click and drag over the text to be deleted. Then press the delete button on your keyboard. The text to be deleted will then be struck through.</p>
Highlight text/ make a comment	<p>Click on the 'Highlight' button on the Commenting tool bar. Click and drag over the text. To make a comment, double click on the highlighted text and simply start typing.</p>	<p>Click on the 'Highlight Text' icon on the Comment tool bar. Click and drag over the text. To make a comment, double click on the highlighted text and simply start typing.</p>
Attach a file	<p>Click on the 'Attach a File' button on the Commenting tool bar. Click on the figure, table or formatted text to be replaced. A window will automatically open allowing you to attach the file. To make a comment, go to 'General' in the 'Properties' window, and then 'Description'. A graphic will appear in the PDF file indicating the insertion of a file.</p>	<p>Click on the 'Attach File' icon on the Comment tool bar. Click on the figure, table or formatted text to be replaced. A window will automatically open allowing you to attach the file. A graphic will appear indicating the insertion of a file.</p>
Leave a note/ comment	<p>Click on the 'Note Tool' button on the Commenting tool bar. Click to set the location of the note on the document and simply start typing. <u>Do not use this feature to make text edits.</u></p>	<p>Click on the 'Add Sticky Note' icon on the Comment tool bar. Click to set the location of the note on the document and simply start typing. <u>Do not use this feature to make text edits.</u></p>

HOW TO...		
Action	Adobe Reader version 7, 8 or 9	Adobe Reader version X
Review	To review your changes, click on the 'Show' button on the Commenting tool bar. Choose 'Show Comments List'. Navigate by clicking on a correction in the list. Alternatively, double click on any mark-up to open the commenting box.	Your changes will appear automatically in a list below the Comment tool bar. Navigate by clicking on a correction in the list. Alternatively, double click on any mark-up to open the commenting box.
Undo/delete change	To undo any changes made, use the right click button on your mouse (for PCs, Ctrl-Click for the Mac). Alternatively click on 'Edit' in the main Adobe menu and then 'Undo'. You can also delete edits using the right click (Ctrl-click on the Mac) and selecting 'Delete'.	To undo any changes made, use the right click button on your mouse (for PCs, Ctrl-Click for the Mac). Alternatively click on 'Edit' in the main Adobe menu and then 'Undo'. You can also delete edits using the right click (Ctrl-click on the Mac) and selecting 'Delete'.

PLEASE DO NOT ATTEMPT TO EDIT THE ARTICLE TEXT ITSELF

SEND YOUR ANNOTATED PDF FILE BACK TO ELSEVIER

Save the annotations to your file and return as an e-mail attachment using the 'reply' button to the original mail. Before returning, please ensure you have answered any questions raised on the Query Form and that you have inserted all corrections: later inclusion of any subsequent corrections cannot be guaranteed.

FURTHER POINTS

- Any (grey) halftones (photographs, micrographs, etc.) are best viewed on screen, for which they are optimized, and your local printer may not be able to output the greys correctly.
- If the PDF files contain colour images, and if you do have a local colour printer available, then it will be likely that you will not be able to correctly reproduce the colours on it, as local variations can occur.
- If you print the PDF file attached, and notice some 'non-standard' output, please check if the problem is also present on screen. If the correct printer driver for your printer is not installed on your PC, the printed output will be distorted.

Our reference: THE 11984

AUTHOR QUERY FORM

Journal: THE Article Number: 11984	Please email or fax your responses and any corrections to: E-mail: Fax:
---	---	---

Dear Author:

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. Note: if you opt to annotate the file with software other than Adobe Reader then please also highlight the appropriate place in the PDF file. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult www.elsevier.com/artworkinstructions.

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the 'AQ' link to go to the location in the proof.

Location in article	Query / Remark: click on the AQ link to go Please insert your reply or correction at the corresponding line in the proof
AQau	Please confirm that given names and surnames have been identified correctly.
AQ1	Ed: Table body does not contain links for †, * and ** in Table 1. Please move †, * and ** to proper locations or delete them and their corresponding table footnotes.
AQ2	Graphics for all figures were supplied as poor quality. Okay as is or supply new figures?

Thank you for your assistance.

ELSEVIER

Available online at www.sciencedirect.com

SciVerse ScienceDirect

Theriogenology

Theriogenology xx (2011) xxx

www.theriojournal.com

Morphometrically-distinct sperm subpopulations defined by a multistep statistical procedure in Ram ejaculates: intra- and interindividual variation

A. Maroto-Morales^a, M. Ramón^b, O. García-Álvarez^b, A.J. Soler^a,
M.R. Fernández-Santos^a, E.R.S. Roldan^c, M. Gomendio, M.D.^c, Pérez- Guzmán^b,
J.J. Garde^{a,d,*}

^a Biology of Reproduction Group, National Wildlife Research Institute (IREC) UCLM-CSIC-JCCM, 02071 Albacete, Spain

^b Regional Center of Animal Selection and Reproduction (CERSYRA) JCCM, 13300 Valdepeñas, Spain

^c Reproductive Ecology and Biology Group, Museo Nacional de Ciencias Naturales (CSIC), 28006 Madrid, Spain

^d Institute for Regional Development (IDR), UCLM, 02071 Albacete, Spain

Received 16 July 2011; received in revised form 27 November 2011; accepted 27 November 2011

Abstract

The existence of sperm subpopulations within the mammalian ejaculate has now been widely recognized. However, to the best of our knowledge, no data exist regarding the existence of sperm morphometric subpopulations within the ovine ejaculate. Computer assisted sperm morphometry analysis (ASMA) data and clustering methods were used in this study to identify sperm-head subpopulations in ram semen. Two experiments were carried out. In Experiment 1, ejaculates from 226 mature rams of the Manchega breed belonging to 36 different herds were used. A minimum of 100 sperm heads were analyzed from each male and eight morphometric characteristics for each individual sperm were recorded. Subpopulation analysis was performed in sequential steps: variable group analysis and correlation analysis to select which morphometric characteristics to use in cluster analyses; nonhierarchical clustering analysis using sperm head length and p2a (also known as roundness) shape factor as initial classificatory variables; and hierarchical clustering analysis to obtain the final number of clusters. The clustering analyses, based on 26 306 individual cells, revealed the existence of four sperm subpopulations (SP1, SP2, SP3 and SP4) with different morphometric characteristics. Significant differences in the proportion of spermatozoa in the SP1 and SP3 were found between rams belonging to different herds. In Experiment 2, the intra- and intermale variability on the distribution of sperm subpopulations was assessed. Three ejaculates from each of 21 rams were collected and the same multistep clustering analysis was performed. For all subpopulations defined, the intermale variability resulted in high values, being the intramale variability much lower. This fact would allow the use of sperm head morphometry to characterize a male and might provide valuable information to assess its fertility. In conclusion, our results show that using computer assisted sperm morphometry analysis and multivariate cluster analyses, four sperm subpopulations with different head phenotype were identified in ram ejaculates.

© 2011 Elsevier Inc. All rights reserved.

Keywords: Sperm morphometry; ASMA; Multistep cluster; Sperm subpopulations; Ram

1. Introduction

The classical approach, considering the whole ejaculate as a homogenous population with a normal statistical distribution, and the use of mean values to

* Corresponding author. Tel: +34 967 599 200+2829; fax: +34 967 599 238.

E-mail address: julian.garde@uclm.es (J.J. Garde).

classify the ejaculates, or to assess the effect of a treatment or a biotechnological procedure is considered incomplete [1–3]. Nowadays, the existence of sperm subpopulations in the mammalian ejaculate is widely recognized [4]. There is evidence that the heterogeneity among these subpopulations may have functional significance [5–7]. Although the origin of these subpopulations and their physiological role are not yet clear [8], it has been found that the ability to undergo capacitation and fertility may vary depending on the subpopulation under consideration [5–9]. Thus, identification of subpopulations has been carried out based on different sperm characteristics, such as biochemical parameters [10–12], functional tests [13–15] or sperm motility descriptors [16–18]. Different statistical procedures to identify and characterize sperm subpopulations have been used [19]. Some of them have been focused on a comparison of different methods to obtain subpopulations, while others have used simple methods providing useful information about the subpopulations identified. These studies have also explored relations between subpopulations identified and sperm quality [20], fertility [21], or their ability to survive cryopreservation [3,22].

Improvements in computer assisted sperm morphometry analysis (ASMA) have also enabled individual sperm to be distinguished in morphometry analysis and subpopulations characterized by morphometric parameters to be identified by multivariate statistical analyses [23–25]. Thus, the existence of sperm-head morphometric subpopulations has been identified in different mammalian species [8,23,25–27]. To our knowledge, little attention has been paid to the study of sperm-head morphometric subpopulations in sheep using ASMA. To date, there has been only one study describing the use of ASMA in the ram [28], and no information is available about the sperm subpopulation structure in fresh ram ejaculates. Previous work [28] has characterized morphometrically the frozen-thawed spermatozoa of this species. Furthermore, this study used a rather small number of animals (i.e., $n = 8$).

Manchega sheep is an autochthonous dairy breed from Spain, which includes a white and a black variety. The white Manchega sheep variety is one of the most important Spanish dairy breeds, widely distributed in the central area of Spain. In this breed, males have not yet been genetically selected for fertility. The average fertility of Manchega rams following artificial insemination at an induced estrus is about 42%, ranging from 8 to 90% [29]. Therefore, males selected for traits, such as milk production are expected to exhibit considerable

diversity in sperm characteristics and in fertility, thus being an excellent model to study the sperm morphometric subpopulation distribution in this species.

Some studies have pointed out the existence of this variability between males according to different sperm characteristics [6,30]. However no study has yet deal the intramale variability in the distribution of the sperm subpopulations. The study of sperm subpopulations distribution could be of utmost biological importance to improve our knowledge about sperm physiology and cryobiology.

Considering this background, the purpose of the present work was mainly to develop a simple multistep statistical procedure to identify, in a large-scale study, sperm-head morphometric subpopulations in an ejaculate based on data gathered with ASMA (Experiment 1), as a basis for future analyses of the relationships between sperm quality, freezability and male fertility. A further aim was to explore if the differences between male variability was affected by the herds of origin of males. Furthermore, we studied the degree of variation in the sperm subpopulations distribution between and within individual sires (21 rams; Experiment 2).

2. Materials and methods

2.1. Animals and reagents

Computer-assisted sperm head morphometry analysis was performed on fresh ejaculates from 226 rams of Manchega sheep breed belonging to 36 herds. Animals were manipulated in accordance with the Spanish Animal Protection Regulation, RD1201/2005, which conforms to European Union Regulation 2010/63. Adult rams were maintained at the Regional Centre of Animal Selection and Reproduction (CERSYRA) located in Valdepeñas, Ciudad Real, (Spain). Ram calves were purchased based on their expected genetic value for milk production. At approximately 3 to 4 mo of age, these rams were transferred from 36 different herds to the AI Centre (CERSYRA), where, after quarantine and training periods of 4 mo, semen was collected. Thus, all males were maintained under same environmental conditions since they were 3 to 4 months old.

All chemicals were of reagent grade and were purchased from Sigma or Merck (both of Madrid, Spain).

2.2. Semen collection

All semen samples were collected with an artificial vagina. The males were all regularly collected (twice a week) in the weeks preceding this study. Semen vol-

ume, sperm concentration and subjective scores of motility (wave motion) were assessed shortly after collection. Volume of each ejaculate was directly measured in graduated tubes. Concentration was estimated using a hemocytometer. Wave motion was scored from 1 to 5 on a wet mount of neat semen at $\times 100$ magnification (values ranged from 0: no movement to 5: strong wave movement). Also, within this interval, aliquots were diluted in phosphate buffered saline (PBS) with bovine serum albumin (5 mg/mL) and used to assess individual sperm motility (0–100%). Only, ejaculates with values of wave motion and individual sperm motility >3 and 80%, respectively, were used.

2.3. Morphometric analysis of sperm heads

Computer-assisted sperm head morphometry analysis was performed on fresh semen as described by Maroto-Morales, et al. [29]. Briefly, semen smears were air-dried and stained using a Hemacolor (Merck) procedure. Stained sperm samples were permanently mounted with a coverslip and dibutyl phthalate xylene (DPX). Stained slides were used to perform ASMA using the morphometry module of a commercially available system (Sperm-Class Analyzer, SCA, Microptic, Barcelona, Spain). The machine was equipped with a Nikon (Labophot-2, Tokyo, Japan) microscope with a $40\times$ bright-field objective and a Sony video camera (CCD AVC-D7CE; Sony Corporation, Tokyo, Japan) connected to a Pentium 950 MHz processor. The illumination source was centered and the intensity of the bulb and the gain and offset of the camera were standardized for all samples. The configuration of the computer system included a PIP-1024 B video digitizer board (Matrox Electronic Systems, Ltd, Quebec, Canada), the sperm image analysis software and a high-resolution assistant monitor Sony Trinitron PVM-1443 MD (Sony Corporation). The array size of the video frame recorder was $512 \times 512 \times 8$ b, digitized images were made up of 262 144 pixels (picture elements) and 256 gray levels. Resolution of images was 0.15 and $0.11 \mu\text{m}$ per pixel in the horizontal and vertical axes, respectively.

The morphometric dimensions for head area (A , μm^2), head perimeter (P , μm), head length (L , μm), head width (W , μm), and four derived parameters of head shape: ellipticity (L/W), $p2a$ ($P [2]/4\pi A$), also known as roundness, elongation ($(L-W)/(L+W)$), and regularity ($\pi LW/4A$), were acquired from 120 to 125 images assuring a minimum of 100 properly measured sperm heads after improperly measured sperm heads were removed from the analysis. The shape feature

roundness ($p2a$) compares the perimeter of an object to its area [31]. This parameter takes a minimum value of one for a perfect circle and is greater than one for more elongated objects. The measurements of each individual sperm head from each ejaculate were saved in an Excel (Microsoft Corporation, Redmond, WA, USA) compatible database by the sperm image analysis software for further analysis.

2.4. Experimental design

2.4.1. Experiment 1: clustering analysis and subpopulation structure

The aim of this experiment was to develop a simple multistep statistical procedure to identify, in a large-scale study, sperm-head morphometric subpopulations within the ram ejaculate. A further aim was to explore the variation in the sperm subpopulations distribution between males belonging to different herds (i.e., origin). For this experiment, ejaculates from 226 mature Manchega rams (*Ovis aries*) belonging to 36 different dairy herds were used.

2.4.2. Experiment 2: study of intra- and inter-ram variability

To provide a more detailed explanation about the male effect on sperm subpopulations, we evaluated the intra- and intermale variability in the frequencies of spermatozoa allocated to each of the four morphometric sperm subpopulations. The objective was to confirm if sperm subpopulation distribution remains constant among the different ejaculates from a male and, therefore, if it would be sufficient to evaluate just one semen sample to assess the morphometric sperm population structure of that male. For this end, three different ejaculates were collected in different days from 21 males in the same season and following a regular regimen (twice a week). A total of 6408 measures of sperm head dimensions were obtained.

2.5. Statistical analysis

All statistical analyses were carried out using the R statistical environment [32]. Where applicable, $P < 0.05$ was considered as statistically significant unless stated otherwise.

A cluster procedure was performed following the multistep process described by Martínez-Pastor, et al. [33]. In a first step, the initial sperm head morphometric parameters to use in the cluster procedure were selected based on the results of a Variable Group Analysis (VGA) and Correlation Analysis among variables (Fig. 1A, Table 1). The objective was to use the lower num-

98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
F149

Fig. 1. Steps followed in the multistep clustering procedure to the identification of the sperm subpopulations. (A) Variable group analysis helps to decide which parameters use as initial classifiers, for this study Length and p2a. (B) Maximum average silhouette width was used in the non-hierarchical clustering as criteria to select the number of clusters which best classify data. For each number of clusters, vertical bars represent the percentage of those clusters. (C) Hubert gamma and the L method were considered in the hierarchical clustering to determine the number of clusters that best classify the data. (D) The initial 11 clusters obtained from the non-hierarchical clustering were finally grouped into 4 clusters in the hierarchical clustering.

ber of variables that allow us to classify spermatozoa based on their sperm head morphometry. The sperm morphometric parameters selected for clustering were the sperm head length and the p2a parameter.

Cluster analysis was divided in two parts: (1) a nonhierarchical cluster analysis using length and p2a as classifiers and, (2) a hierarchical cluster analysis per-

formed on the clusters obtained from the previous analysis. Non-hierarchical methods are often used with large data sets before trying a hierarchical analysis to reduce the data to a few initial clusters and then pass them to the hierarchical procedure. Nonhierarchical analysis was carried out using the Clara function on the CLUSTER package. The algorithm is based on the

Table 1

Correlation analysis among ram sperm head morphometric parameters. Pearson's correlation values are presented on lower diagonal and significance level are on upper diagonal. †, **, ***

	Length	Width	Area	Perimeter	p2a	Ellipticity	Elongation	Regularity
Length		a	a	a	a	a	a	a
Width	0.373		a	a	a	a	a	a
Area	0.667	0.823		a	a	a	a	a
Perimeter	0.410	0.272	0.438		a	a	a	a
p2a	0.042	-0.200	-0.127	0.832		a	a	NS
Ellipticity	0.424	-0.676	-0.273	0.070	0.242		a	a
Elongation	0.425	-0.678	-0.274	0.062	0.233	0.997		a
Regularity	0.474	0.351	0.077	0.039	-0.002	0.027	0.029	

† NS: non-significant; * $P \leq 0.05$; ** $P < 0.01$; a $P < 0.001$.

Numbers in bold and italic represent the correlations among sperm parameters initially considered for the characterization of sperm subpopulations.

search for k representative medoids among the observations of the dataset and subsequent construction of k clusters which minimize the sum of the dissimilarities of the observations to their closest representative medoid. Euclidean distances, based on root sum-of-squares of differences, were used for calculating dissimilarities between observations. Previous to the nonhierarchical procedure, a standardization of data was performed to avoid the negative effects of considering classification variables measured in different scales. The optimal numbers of clusters to keep and use in the following hierarchical analysis was set by using the Silhouette Average Width (SAW) criterion [34]. The Silhouette validation technique calculates the silhouette width for each sample, with values ranging from -1 to 1 . If silhouette value is close to one, it means that sample is "well-clustered" and it was assigned to a very appropriate cluster. If silhouette value is about zero, it means that sample could be assigned to another close cluster as well, and the sample lies equally far away from both clusters. If silhouette value is close to -1 , it means that sample is "misclassified" and is merely somewhere in between the clusters. The overall average silhouette width is the average of each silhouette width for all objects in the whole data set. The largest overall average silhouette indicates the best clustering (number of cluster). Therefore, the number of clusters with maximum average silhouette width is taken as the optimal number of the clusters. For this work, nonhierarchical cluster analysis was carried out considering from 4 to 20 possible clusters. The highest value of SAW was obtained for 11 clusters (Fig. 1B). This clustering procedure allowed us to detect and remove outliers as well because of its sensitivity to them.

Once the nonhierarchical analysis was carried out and 11 initial clusters were obtained, a hierarchical

procedure was performed using the HCLUST function on STATS package. Distance matrix was computed by using the *Euclidean* distance measure and the *Ward's* minimum variance method was set to classify the data. A dendrogram plot showing the clustering process was obtained (Fig. 1C). To determine the final number of clusters two criteria were considered: the *Hubert Γ coefficient* [35] and the *L Materials and methods* [36]. The *Hubert Γ* measures the level of similarity of cluster structure so the higher the value, the better the cluster structure is. This coefficient has been used previously in cluster analyses of spermatozoa [17]. The *L Method* was used because of its simplicity. This method determines the optimal number of clusters based on graphical methods by finding the "knee", that is, the point of maximum curvature in an evaluation graph that is created by the clustering algorithm. The evaluation graph is a two-dimensional plot where the x axis is the number of clusters, and the y axis can be any evaluation metric, such as distance, similarity, error, or quality. In that evaluation plot we usually observed two differentiated regions: one on the right and the other one on the left of the knee and they are often approximately linear. If a line is fitted to the right side and another line is fitted to the left side, then the area between the two lines will be in the same region as the knee. The value of the x axis at the knee can then be used as the number of clusters to return. Following both criteria the optimal number of clusters and therefore the final number of subpopulations was set to four (Fig. 1D).

Once the sperm subpopulations were identified and characterized, different regression analyses were performed to evaluate the subpopulations distribution among herds of origin and to study the intra- and intermale variability. For the latter, a mixed effects model was carried out considering the male as a random effect, and the coefficients of variation (the ratio of

the standard deviation to the mean and expressed as percentage) between males (CVb) and within a male (between ejaculates) (CVw) were also computed as a direct measure of the variability of subpopulation distribution.

3. Results

3.1. Experiment 1: clustering analysis and subpopulation structure

3.1.1. Selection of initial variables and cluster analysis

The sperm head length and the p2a shape factor were chosen as initial variables for clustering based on the results obtained from a VGA and Correlation Analysis. VGA dendrogram (Fig. 1A) showed how variables were grouped into three main groups: head shape parameters tended to group first, followed by head length and head width and finally, the head area and the head perimeter. To classify spermatozoa based on their sperm head morphometry a linear combinations of variables on each group (principal components analysis) could be used or one parameter of each group can be selected. We decided to use raw variables, so we first chose length, area and p2a as possible candidate variables. Correlation between length and p2a resulted close to 0 (Table 1), indicating that both variables are unrelated. Length and area are highly correlated ($r = 0.67$), so we decided to use only sperm head length since its correlations with p2a was lower.

Non-hierarchical cluster analysis using length and p2a was carried out and several 11 clusters were kept based on the *Silhouette Average Width* values (Fig. 1B). The distribution of data on these clusters was considered appropriated (see bars on Fig. 1B), since none of the clusters presented a short number of observations indicating possible outliers (group size ranged from 451 to 4225 observations, with an average size of 2300 observations). The resulting clusters from the non-hierarchical analysis were grouped into four clusters (Fig. 1D) after applying the hierarchical procedure and studying the *Hubert Γ* coefficient and the *L Materials and methods* plot (Fig. 1C).

3.1.2. Sperm subpopulations characteristics

Four sperm subpopulations (clusters) were identified after the cluster analysis of the 26 306 individual spermatozoa (observations). As it can be observed, subpopulations were clearly differentiated by head sperm length and p2a parameters (Table 2). Group size ranged from 451 (SP4) to 12 544 (SP2) observations. Sum-

Table 2

Summary of morphometric characteristics for each sperm subpopulation obtained on Manchega rams. Data are expressed as mean $\pm$ SD.

Subpopulation	n	Length	p2a	Width	Area	Perimeter	Ellipticity	Elongation	Regularity
1	5281	9.57 $\pm$ 0.26 ^a	1.58 $\pm$ 0.22 ^a	4.99 $\pm$ 0.35 ^a	37.96 $\pm$ 3.05 ^a	27.48 $\pm$ 1.96 ^a	1.92 $\pm$ 0.13 ^a	0.31 $\pm$ 0.03 ^a	0.98 $\pm$ 0.04 ^a
2	12,544	8.95 $\pm$ 0.21 ^b	1.70 $\pm$ 0.23 ^b	4.78 $\pm$ 0.29 ^b	35.15 $\pm$ 2.45 ^b	27.03 $\pm$ 1.92 ^b	1.87 $\pm$ 0.12 ^b	0.30 $\pm$ 0.03 ^b	0.96 $\pm$ 0.04 ^b
3	8030	8.40 $\pm$ 0.27 ^c	1.63 $\pm$ 0.20 ^c	4.66 $\pm$ 0.28 ^c	32.68 $\pm$ 2.34 ^c	25.58 $\pm$ 1.72 ^c	1.79 $\pm$ 0.11 ^c	0.28 $\pm$ 0.03 ^c	0.93 $\pm$ 0.04 ^c
4	451	9.64 $\pm$ 0.43 ^d	2.48 $\pm$ 0.24 ^d	4.71 $\pm$ 0.41 ^d	35.23 $\pm$ 3.44 ^b	32.98 $\pm$ 1.73 ^d	1.98 $\pm$ 0.18 ^d	0.33 $\pm$ 0.04 ^d	0.97 $\pm$ 0.05 ^d

^a Values within a column with different superscripts differ ($P < 0.001$).

^d Values within a column with different superscripts differ ($P < 0.001$).

Table 3

Sperm head morphometry subpopulations distribution in males of Manchega sheep breed. Data are means $\pm$ standard errors. All sperm subpopulations were significantly different ($P < 0.001$) between males.

	Sperm subpopulations ^a			
	SP1	SP2	SP3	SP4
Means $\pm$ SEM	29.26 $\pm$ 1.88	40.73 $\pm$ 2.52	23.74 $\pm$ 1.98	6.28 $\pm$ 0.81
Range	1.43–60.00	7.84–79.57	1.96–67.65	0.00–23.71

^a SP1: long and wide sperm head. SP2: small and sharp sperm head. SP3: small and wide sperm head. SP4: long and sharp sperm head.

mary statistics of these subpopulations are presented in Table 2, and the characteristics of each sperm subpopulation based on sperm head length and p2a parameters used on cluster analysis, are the following:

Subpopulation 1 (SP1): This subpopulation included those spermatozoa whose sperm head is long (average length of $9.6 \pm 0.3 \mu\text{m}$) and wide (average p2a of 1.6 ± 0.2). The spermatozoa belonging to this subpopulation are the biggest. This subpopulation was characterized by spermatozoa with the smallest values of p2a. Thus, this is the subpopulation with spermatozoa with the shape of the head more similar to a circle. The $19.3 \pm 14.4\%$ (mean $\pm$ standard deviation) of the spermatozoa in the data set were assigned to this subpopulation.

Subpopulation 2 (SP2): This subpopulation included those spermatozoa whose sperm head was short (length = $8.95 \pm 0.21 \mu\text{m}$), but with high values of p2a (1.70 ± 0.23). This subpopulation contained the largest number of spermatozoa ($47.9 \pm 10.5\%$).

Subpopulation 3 (SP3): This subpopulation was characterized by spermatozoa with the shortest sperm head dimensions (average length of $8.4 \pm 0.3 \mu\text{m}$; average p2a of 1.6 ± 0.2). Spermatozoa belonging to this subpopulation represented the $31.0 \pm 17.2\%$ of the total data.

Subpopulation 4 (SP4): This subpopulation included spermatozoa whose sperm head length (9.6 ± 0.4), and p2a (2.5 ± 0.2) were the highest. Thus, this subpopulation included the spermatozoa with the sharpest (elongated) head. This was the smallest subpopulation, with an average percentage per male of 1.8 ± 3.5 .

3.1.3. Effects of herd of origin of males

Once the spermatozoa were classified by a cluster analysis, the distribution of spermatozoa within each subpopulation was calculated for each male and for each herd of origin of males considered in this experiment. Average percentages per herd were 17.7 ± 8.2 for SP1, 47.2 ± 5.4 for SP2, 33.2 ± 11.2 for SP3; and

1.8 ± 2.0 for SP4. We observed an important variation in the percentages of subpopulations SP1 and SP3, being the herd effect significant in both subpopulations ($P < 0.05$). Regarding with the SP4, differences between herds were lower than for the SP1 and SP3, with a non-significant herd effect ($P > 0.05$). However, the intraherd variability was lower for the SP2 than the others.

3.2. Experiment 2: study of intra- and inter-ram variability

The effect of male in the percentage of the different morphometrically sperm subpopulations is presented on Table 3. All males included in this experiment showed great differences in the distribution of the four subpopulations, being the individual effect of male statistically significant ($P < 0.001$). When comparing the intra- and intermale variability, the latter was always larger with CV_w of 10 to 20% and CV_b of 50 to 70%, reaching even the 100% in the particular case of SP4 (Table 4). The intra- and intermale variability comparison is better observed on Fig. 2, especially for the SP2. Thus, the SP2 is the subpopulations that showed most consistent rates.

4. Discussion

A previous study of our group utilizing ASMA has already established the sperm-head dimensions for ram

Table 4

Coefficients of variation between males and within males (between ejaculates of the same male).

CV %	Sperm subpopulations ^a			
	SP1	SP2	SP3	SP4
Between-male	50.73	49.12	65.98	97.81
Within-male ^b	16.76	12.80	20.33	60.97

^a SP1: long and wide sperm head. SP2: small and sharp sperm head. SP3: small and wide sperm head. SP4: long and sharp sperm head.

^b Average of individual CV within each male.

Fig. 2. Variability on the morphometrically sperm subpopulations between and within (between ejaculates) males. Means (open circles) and SEM (whiskers) and represented. (A) long and wide sperm head (SP1). (B) Small and sharp sperm head (SP2). (C) Small and wide sperm head (SP3). (D) long and sharp sperm head (SP4).

spermatozoa and has revealed that ram ejaculates are very heterogeneous at least in their cell morphometric characteristics [29]. In this study, we have presented a simple method that combines ASMA techniques and a multivariate analysis procedure using commonly used statistical tools to identify, in a large-scale study, sperm

head morphometric subpopulations. The statistical procedure hereby used is simple and appears useful to detect sperm subpopulations. This procedure is a different approach from that used in previous studies. Large number of studies are available on the characterization of sperm subpopulations. A recent review on

the statistical procedures commonly used up to date in the characterization of sperm subpopulations has been provided by Martínez-Pastor, et al. [19].

One of the main objectives of this study has been to provide a detailed description of the procedures that we have used to obtaining the subpopulations. The first step in our clustering procedure has been to determine which sperm head morphometry characteristics use to classify the spermatozoa into different groups. Some authors propose the use of a principal component analysis (PCA) as a previous step before cluster analysis [28,33] to obtain a small number of variables that retain as much of the information in the original variables as possible. In this study, we have preferred to use primary variables rather than principal components since these variables allow us to keep all the variability of data and we have reasoned that the results obtained from a cluster analysis with primary variables would be clearer. From the results of the Variable Analysis Group (VAG) and the Correlation Analysis we have chose the sperm head long and the shape factor p2a as the initial classifiers.

By using ASMA methodology and multivariate cluster analyses, four sperm subpopulations with different morphometric characteristics have been identified in ram ejaculates. These subpopulations have been characterized by different values of sperm-head dimensions and shape factors. These data confirm our idea that ovine ejaculates are not homogeneous samples [29]. Our results agree with those found in other mammalian species [21,26,27]. Regarding to the ram, Martí, et al. [28] have reported different number of sperm subpopulations for younger (3 subpopulations) and older (4 subpopulations) ram using frozen-thawed sperm samples. These authors have performed a PCA previously to the clustering analysis and the subpopulations have been identified by using other sperm morphometric parameters as classifiers. Their results differ from those obtained in our work, mainly because of the effect of cryopreservation, since it has been reported that sperm head size is significantly smaller in cryopreserved spermatozoa compared to fresh-extended sperm cells [26,27,37].

An important finding of our large-scale study has been the existence of considerable variation between individual males in the percentages of spermatozoa within each subpopulation, being the percentages of all subpopulations significantly different between males. In our study, rams were not preselected for fertility or any sperm characteristics. Probably, if male selection had been carried out for fertility, we would expect to

observe a less profound variation in sperm phenotype (such as morphometry) between males. This fact did not happen in our study since males are selected only for milk production, exhibiting a great diversity in sperm head morphometric dimensions and, therefore, in their subpopulation structure. This fact gives more importance to the choice of Manchega ram as animal model.

But possibly, the most important finding of this study has been the relation between inter- and the intramale variability in the distribution of the morphometrically sperm subpopulations. Our results have shown for the percentage of the four sperm subpopulations higher CV_b than CV_w . The most suitable parameters form use in the characterization of individual males should be those characterized by relatively low CV_w and relatively high CV_b [38,39]. So, low intramale variability observed in this study is a good indicator that the sperm subpopulation structure based on the spermatozoa head morphometry is quite constant within a male. Thus, this technique could be very useful for studying properly the relationships between sperm morphometry with several reproductive performances or to evaluate the effects caused by the sperm handling, such as cryopreservation procedure.

Our findings have shown that the contributions of subpopulations SP1 (big and round) and SP2 (small and elongated) could be useful to characterize individuals because of their low CV_w . Some authors have obtained higher fertility rates in stallion when the proportion of spermatozoa with small head has increased in the samples [23]. Similarly, the relationship between sperm head dimensions in fresh ejaculates and sperm freezability has been previously supported for boar spermatozoa [6,37]. Peña, et al. [37] have reported in boar that the percentage of spermatozoa in the fresh ejaculates with the smallest head dimensions was positively correlated with the sperm quality after thawing. So, applying techniques, such as cluster analysis to ASMA data we may be able to further identify those spermatozoa with adaptive differences, such as increased fertilizing ability or resistance to cryoinjury, thus promoting morphology analysis as a valuable diagnostic tool.

The differences in sperm morphometry between males have been widely reported, but our understanding of the causal factors that generate such differences is still poor. Genetically determined variation in sperm morphology has been recognized for some time and was demonstrated clearly in the observation of phenotypic differences between sperm

of different strains of mice [40]. It has been suggested that variation in sperm morphology is originated during spermatogenesis when genotypic effects influence sperm structure [6,41]. Sperm phenotype appears to be controlled by genes transcribed in the premeiotic phase of development (diploid genome) [42]. Clear examples of sperm development and morphology under strict genetic control have been demonstrated in studies linking inbreeding coefficients and poor ejaculate quality in endangered ungulates [41]. Furthermore, studies in fish species with high sperm competition levels have showed that sperm quality varies between males [43]. Therefore, it is reasonable to suggest that the between-male differences reported here are under genetic control.

Once we have established that between-rams differences in sperm subpopulations exist, it is important to determine if differences observed in the percentages of spermatozoa belonging to each subpopulation between rams could be due to origin (i.e., herd). Our findings indicate that there are differences in the distribution of spermatozoa within each subpopulation between rams belonging to different herds (i.e., origin). Given that, in this sample of males belonging to 36 populations, the environmental factors were common to all individuals since they were 3 to 4 months old, the effect of the herd of origin on sperm morphometry could suggest a genetic effect. The combination of an individual and herd effects builds strong support for the view that variation in ram sperm head morphometry (phenotype) exists and may be genetically inherited.

In conclusion, our results show that using ASMA and multivariate cluster analyses, four sperm subpopulations with different head phenotype were identified in ram ejaculates. Considerable variations between rams in the percentages of spermatozoa allocated to each subpopulation were observed and a further study allowed us to reveal that these differences were significant for all subpopulations. The herd of origin of males influenced subpopulation distribution. Considering that environmental factors were common to all individuals since they were 3 months old, the effect of the herd of origin on sperm morphometry suggests a genetic control. Further research using ASMA would be useful to identify possible effects of cryopreservation on ram spermatozoa distribution in morphometrically subpopulations and the relationships between morphometric sperm subpopulations distribution and other sperm functional characteristics.

Acknowledgments

This work has been supported by the Education and Science Council of Junta de Comunidades de Castilla-La Mancha (PBC-05-008). A Maroto-Morales and O García-Álvarez were recipients of scholarships from Junta de Comunidades de Castilla-La Mancha and Inia, respectively.

References

- [1] Mortimer ST. A critical review of the physiological importance and analysis of sperm movement in mammals. *Hum Reprod Update* 1997;3:403–39.
- [2] Mortimer ST. CASA-practical aspects. *J Androl* 2000;2000: 514–24.
- [3] Ortega-Ferrusola C, Macías García B, Suárez Rama V, Gallardo-Bolaños JM, González-Fernández L, Tapia JA, et al. Identification of sperm subpopulations in stallion ejaculates: changes after cryopreservation and comparison with traditional statistics. *Reprod Domest Anim* 2009;44:419–>23.
- [4] Holt WV, Van Look KJ. Concepts in sperm heterogeneity, sperm selection and sperm competition as biological foundations for laboratory tests of semen quality. *Reproduction* 2004; 127:527–35.
- [5] Holt WV. Can we predict fertility rates? Making sense of sperm motility. *Reprod Dom Anim* 1996;31:17–24.
- [6] Thurston LM, Watson PF, Mileham AJ, Holt WV. Morphologically distinct sperm subpopulations defined by Fourier shape descriptors in fresh ejaculates correlate with variation in boar semen quality following cryopreservation. *J Androl* 2001;22: 382–94.
- [7] Núñez-Martínez I, Moran JM, Peña FJ. Two-step cluster procedure after principal component analysis identifies sperm subpopulations in canine ejaculates and its relation to cryoresistance. *J Androl* 2006a;27:596–603.
- [8] Thurston LM, Watson PF, Holt WV. Sources of variation in the morphological characteristics of sperm subpopulations assessed objectively by a novel automated sperm morphology analysis system. *J Reprod Fertil* 1999;117:271–80.
- [9] Harrison RA. Capacitation mechanisms, and the role of capacitation as seen in eutherian mammals. *Reprod Fertil Dev* 1996; 8:581–94.
- [10] Calamera J, Buffone M, Ollero M, Alvarez J, Doncel GF. Superoxide dismutase content and fatty acid composition in subsets of human spermatozoa from normozoospermic, asthenozoospermic, and polyzoospermic semen samples. *Mol Reprod Dev* 2003;66:422–30.
- [11] Paasch U, Grunewald S, Fitzl G, Glander HJ. Deterioration of plasma membrane is associated with activated caspases in human spermatozoa. *J Androl* 2003;24:246–52.
- [12] Buffone MG, Doncel GF, Marín Briggiler CI, Vazquez-Levin MH, Calamera JC. Human sperm subpopulations: relationship between functional quality and protein tyrosine phosphorylation. *Hum Reprod* 2004;19:139–46.
- [13] Holt C, Holt WV, Moore HD, Reed HC, Curnock RM. Objectively measured boar sperm motility parameters correlate with the outcomes of on-farm inseminations: results of two fertility trials. *J Androl* 1997;18:312–23.

- [14] Petrunkina AM, Töpfer-Petersen E. Heterogeneous osmotic behaviour in boar sperm populations and its relevance for detection of changes in plasma membrane. *Reprod Fertil Dev* 2000; 12:297–305.
- [15] Pérez-Llano B, Yenes-García P, García-Casado P. Four subpopulations of boar spermatozoa defined according to their response to the short hypoosmotic swelling test and acrosome status during incubation at 37 degrees C. *Theriogenology* 2003; 60:1401–7.
- [16] Nuñez-Martínez I, Moran JM, Peña FJ. A three-step statistical procedure to identify sperm kinematic subpopulations in canine ejaculates: changes after cryopreservation. *Reprod Dom Anim* 2006b;41:408–15.
- [17] Martínez-Pastor F, Cabrita E, Soares F, Anel L, Dinis MT. Multivariate cluster analysis to study motility activation of *Solea senegalensis* spermatozoa: A model for marine teleosts. *Reproduction* 2008;135:449–59.
- [18] Muñio R, Peña AI, Rodríguez A, Tamargo C, Hidalgo CO. Effects of cryopreservation on the motile sperm subpopulations in semen from asturiana de los Valles bulls. *Theriogenology* 2009;72:860–8.
- [19] Martínez-Pastor F, Tizado EJ, Garde JJ, Anel L, de Paz P. Statistical series: opportunities and challenges of sperm motility subpopulation analysis. *Theriogenology* 2011;75:783–95.
- [20] Saravia F, Núñez-Martínez I, Morán JM, Soler C, Muriel A, Rodríguez-Martínez H, et al. Differences in boar sperm head shape and dimensions recorded by computer-assisted sperm morphometry are not related to chromatin integrity. *Theriogenology* 2007;68:196–203.
- [21] Quintero-Moreno A, Miró J, Teresa Rigau A, Rodríguez-Gil JE. Identification of sperm subpopulations with specific motility characteristics in stallion ejaculates. *Theriogenology* 2003;59: 1973–90.
- [22] Beirão J, Cabrita E, Pérez-Cerezales S, Martínez-Páramo S, Herráez MP. Effect of cryopreservation on fish sperm subpopulations. *Cryobiology* 2011;62:22–31.
- [23] Gravance CG, Liu IK, Davis RO, Hughes JP, Casey PJ. Quantification of normal head morphometry of stallion spermatozoa. *J Reprod Fertil* 1996;108:41–6.
- [24] Estes MC, Fernandez-Santos MR, Soler AJ, Garde JJ. Head dimensions of cryopreserved red deer spermatozoa are affected by thawing procedure. *Cryo Lett* 2003;24:261–8.
- [25] Alvarez M, García-Macías V, Martínez-Pastor F, Martínez F, Borragán S, Mata M, et al. Effects of cryopreservation on head morphometry and its relation with chromatin status in brown bear (*Ursus arctos*) spermatozoa. *Theriogenology* 2008;70: 1498–506.
- [26] Rubio-Guillén J, González D, Garde JJ, Estes MC, Fernández-Santos MR, Rodríguez-Gil JE, et al. Effects of cryopreservation on bull spermatozoa distribution in morphometrically distinct subpopulations. *Reprod Dom Anim* 2007;42:354–7.
- [27] Estes MC, Fernández-Santos MR, Soler AJ, Montoro V, Martínez-Pastor F, Garde JJ. Identification of sperm-head subpopulations in Iberian red deer epididymal sperm samples. *Reprod Dom Anim* 2009;44:206–11.
- [28] Martí JJ, Aparicio IM, García-Herreros M. Sperm morphometric subpopulations are differentially distributed in rams with different maturity age in cryopreserved ejaculates. *Theriogenology* 2011;76:97–109.
- [29] Maroto-Morales A, Ramón M, García-Alvarez O, Soler AJ, Estes MC, Martínez-Pastor F, et al. Characterization of ram (*Ovis aries*) sperm head morphometry using the sperm-Class analyzer. *Theriogenology* 2010;73:437–48.
- [30] Alvarez C, Castilla JA, Martínez L, Ramírez JP, Vergara F, Gaforio JJ. Biological variation of seminal parameters in healthy subjects. *Hum Reprod* 2003;18:2082–8.
- [31] Sailer BL, Jost LK, Evenson DP. Bull sperm head morphometry related to abnormal chromatin structure and fertility. *Cytometry* 1996;24:167–73.
- [32] R Development Core Team. A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2009.
- [33] Martínez-Pastor F, García-Macias V, Alvarez M, Herraez P, Anel L, de Paz P. Sperm subpopulations in Iberian Red deer epididymal sperm and their changes through the cryopreservation process. *Biol Reprod* 2005;72:316–27.
- [34] Silhouettes RPJ. A graphical aid to the interpretation and validation of cluster analysis. *J Comput Appl Math* 1987;20:53–65.
- [35] Haldiki M, Batistakis Y, Vazirgiannis M. Cluster validity methods, SIGMOD. Record;2002:405.
- [36] Salvador S, Chan P. Determining the number of clusters/segments in hierarchical clustering/segmentation algorithms, in IC-TAI. IEEE Computer Society; 2004, p. 576–84.
- [37] Peña FJ, Saravia F, García-Herreros M, Núñez-Martínez I, Tapia JA, Johannisson A, et al. Identification of sperm morphometric subpopulations in two different portions of the boar ejaculate and its relation to postthaw quality. *J Androl* 2005;26: 716–23.
- [38] Gago C, Pérez-Sánchez F, Yeung CH, Tablado L, Cooper TG, Soler C. Morphological characterization of ejaculated cynomolgus monkey (*Macaca fascicularis*) sperm. *Am J Primatol* 1999; 47:105–15.
- [39] Phetudomsinsuk K, Sirinarumitr K, Laikul A, Pinyopummin A. Morphology and head morphometric characters of sperm in Thai native crossbred stallions. *Acta Vet Scand* 2008;50:41.
- [40] Beatty RA, Sharma KN. Genetic of gametes II. Strain differences in spermatozoa from 8 inbred strains of mice. *Proc Biol Sci* 1960;1960:25–53.
- [41] Roldan ER, Cassinello J, Abaigar T, Gomendio M. Inbreeding, fluctuating asymmetry, and ejaculate quality in an endangered ungulate. *Proc Biol Sci* 1998;265:243–8.
- [42] Morrow EH, Gage MJG. Consistent significant variation between individual males in spermatozoal morphometry. *J Zool* 2001;254:147–53.
- [43] Fitzpatrick JL, Desjardins JK, Milligan N, Montgomerie R, Balshine S. Reproductive-tactic-specific variation in sperm swimming speeds in a shell-brooding cichlid. *Biol Reprod* 2007;77:280–4.