

Review

Tick–Pathogen Interactions: The Metabolic Perspective

Alejandro Cabezas-Cruz,^{1,*} Pedro Espinosa,² Pilar Alberdi,² and José de la Fuente^{2,3,*}

The first tick genome published in 2016 provided an invaluable tool for studying the molecular basis of tick–pathogen interactions. Metabolism is a key element in host–pathogen interactions. However, our knowledge of tick–pathogen metabolic interactions is very limited. Recently, a systems biology approach, using omics datasets, has revealed that tick-borne pathogen infection induces transcriptional reprogramming affecting several metabolic pathways in ticks, facilitating infection, multiplication, and transmission. Results suggest that the response of tick cells to tick-borne pathogens is associated with tolerance to infection. Here we review our current understanding of the modulation of tick metabolism by tick-borne pathogens, with a focus on the model intracellular bacterium *Anaplasma phagocytophilum*.

Tick–Pathogen Metabolic Interactions at the Center of the Infectious Storm

Metabolism is key to cellular function [1,2]. In consequence, all major molecular pathways in living cells are interconnected to, and regulated by, the availability and levels of metabolites [1,2]. Recent studies in mammalian cells showed that interactions between intracellular bacterial pathogens and the host cells can lead to physiological changes in both interacting members [3]. The metabolic adaptations in mammalian cell–pathogen systems promote proliferation or elimination of the pathogen within the host cells [3]. Tick–pathogen interactions, however, involve a more complex array of outcomes which include conflict and cooperation that ultimately benefit both ticks and pathogens [4]. This is not surprising considering that, in vector-borne pathogen systems, the survival of the vector is essential for the completion of the life cycle of the pathogen. In addition, tick-borne pathogen infection increases tick performance in challenging environmental conditions [5,6]. Therefore, in some cases, vector tolerance to pathogen infection is an advantageous life trait in arthropod vectors [7].

Modulation of tick metabolism by tick-borne pathogens is a result of coevolution and adaption to a considerable number of tick and reservoir host species [4,8,9]. The *Ixodes scapularis* genome, the first for a medically important chelicerate, is the only tick genome available so far [10]. The sequencing of the *I. scapularis* genome was an essential step towards the understanding of the molecular processes that support the parasitic lifestyle of the tick and its success as a vector of multiple pathogens, including bacteria, viruses, protozoa, and helminths, which constitute a growing burden for human and animal health worldwide [11]. This review focuses on recent research that provided insights on the finely tuned metabolic changes that the model pathogen *A. phagocytophilum* induces in its tick vector *I. scapularis*. For comparative purposes, tick metabolic changes induced by other pathogens, including mainly *Borrelia* spp., flaviviruses, and fungi, will be also included (Table 1). Results suggest that tick-borne pathogens use tick metabolism as a hub to modulate tick physiology and tick cell processes, including immunity and apoptosis among others.

Highlights

The first tick genome sequence has greatly improved our understanding of the molecular basis of tick metabolism and its interaction with tick-borne pathogens.

Tick-borne pathogens use tick metabolism as a hub to modulate tick physiology and tick cell processes, including immunity and apoptosis among others.

Anaplasma phagocytophilum rewires the tick metabolic network and changes the relative importance of some pathways, including gluconeogenesis and glyceroneogenesis.

Fatty acids play important roles as energy sources for many pathogens and ticks, which have to endure long periods of time off the host between feedings.

¹UMR BIPAR, INRA, Ecole Nationale Vétérinaire d'Alfort, ANSES, Université Paris-Est, Maisons-Alfort, France

²SaBio, Instituto de Investigación en Recursos Cinegéticos IREC (CSIC-UCLM-JCCM), Ciudad Real, Spain

³Department of Veterinary Pathobiology, Center for Veterinary Health Sciences, Oklahoma State University, Stillwater, OK, USA

*Correspondence: cabezasalejandrocruz@gmail.com (A. Cabezas-Cruz) and jose_delafuente@yahoo.com (J. de la Fuente).

Nutritional Immunity and Iron Metabolism in Ticks

Iron is essential for most organisms because it serves as an electron donor and acceptor in various metabolic processes. Therefore, iron holds a central position in host–pathogen interactions [12]. While hosts evolved mechanisms to deprive pathogens of iron, pathogens coevolved mechanisms to acquire the host's iron. Host mechanisms involved in limiting the access of pathogens to iron were termed 'nutritional immunity' [12]. During feeding, ticks are exposed to the large amounts of iron present in the blood meal. Excess iron can react with H_2O_2 , yielding hydroxyl radical, a potent biological oxidant [12]. Therefore, a balanced regulation of iron levels is essential for ticks [13], and potentially, pathogen survival. Two ferritin (FER) proteins, FER1 and FER2, are the primary iron storage and transporter, respectively, in ticks [13]. In addition to controlling the production of reactive oxygen species (ROS) from iron, as in other model organisms [12,13], ferritins may also contribute to nutritional immunity in ticks by depriving pathogens of their essential supply of iron. Interestingly, *Escherichia coli* infection induces the expression of a *fer* gene in *Dermacentor variabilis*, and silencing of *fer* genes in *Haemaphysalis longicornis* decreases tick survival following a challenge with *E. coli* [13].

The tick-borne pathogen *Ehrlichia ruminantium* senses iron levels in mammalian cells and responds to iron starvation by upregulating virulence genes such as those involved in the type IV secretion system (T4SS), which, in turn, may participate in iron acquisition by the pathogen [14]. A similar mechanism may be at play during pathogen infection in ticks. An iron-binding protein expressed in the infective stage of *A. phagocytophilum* [15], and hypothesized to be secreted by the T4SS, was found in tick salivary glands and tick cells infected with these bacteria [16]. Bacterial iron-binding proteins may sequester iron to overcome nutritional immunity in ticks. An interesting mechanism used to escape nutritional immunity is that of *Borrelia burgdorferi* which requires no, or very low amounts of, iron to survive [17]. In addition, recent research provides evidence that *A. phagocytophilum* modulates tick heme metabolism. Host hemoglobin is cleaved to heme by three sequential reactions involving several hemoglobinolytic enzymes in ticks [18]. Due to the high redox potential of heme, this molecule is toxic and, in excess, can be bacteriostatic [19]. Quantitative proteomics analysis revealed that *A. phagocytophilum* modulates the levels of hemoglobinolytic enzymes, affecting the levels of host hemoglobin in tick midguts and salivary glands [18]. This finding suggests that *A. phagocytophilum* infection may regulate the levels of heme in a tissue-specific manner with potential effects for pathogen and vector survival.

Tick Hemocyte Metabolism in Response to Infection by Tick-borne Pathogens

Despite mammalian macrophages possess specialized effector functions to eliminate invading microorganisms, many intracellular bacteria infect and replicate inside these immune cells [20]. Globally, macrophages can be divided into inflammatory (M1) and anti-inflammatory (M2) macrophages [20]. Both types of macrophage, M1 and M2, can be infected by intracellular bacteria; however, the metabolic profiles of these macrophages in response to pathogenic stimulus are substantially different [20]. After stimulation with lipopolysaccharide (LPS), M1 macrophages take up more glucose via **glucose transporters (GLUTs)**, see [Glossary](#) and increase the glycolytic flux and the production of lactate, a process named **aerobic glycolysis**. In contrast, M2 macrophages have a metabolic profile comparable to that of unstimulated cells, but with higher **tricarboxylic acid cycle (TCA cycle)** and oxidative activity [20].

Cellular immunity in ticks is mostly mediated by hemocytes, which are involved in the phagocytosis of different microbes [21]. After establishing a primary infection in tick midguts, *A. phagocytophilum* migrates to the salivary glands [11]. A secreted tick protein, P11, enables *A.*

Glossary

Acetyl-CoA: a central metabolic intermediate synthesized in the mitochondria. Acetyl-CoA is the molecule through which glycolytic pyruvate enters the TCA cycle to be oxidized for energy production. It can be derived from carbohydrates, fatty acids, and amino acids.

Aerobic glycolysis: a term frequently used to name a metabolic profile similar to that of the Warburg effect in which a cell, even under normoxic conditions, generates a high amount of energy by using the glycolytic pathway, notably with a high production of lactate.

β -Oxidation: the major pathway for the degradation of fatty acids; it is essential for producing energy from the generation of acetyl-CoA, NADH, and $FADH_2$ when glucose supply is limiting.

Gluconeogenesis: a metabolic pathway that, during a longer-term fast or starvation, results in *de novo* glucose synthesis from noncarbohydrate carbon substrates, mainly lactate and glycerol.

Glucose transporters (GLUTs): a wide group of membrane proteins that transport glucose across the cell membrane. The GLUT family is divided into three classes of which class I member GLUT1 is transcriptionally regulated by HIF-1 α .

Glyceroneogenesis: a metabolic pathway that is activated during glucose deprivation to synthesize glycerol-3-phosphate, a building block of triglyceride and fatty acids. In this pathway, glycerol-3-phosphate is synthesized from pyruvate, alanine, glutamine, and TCA cycle intermediates.

Glycolysis: a major catabolic pathway in which one molecule of glucose is transformed into two molecules of pyruvate, in the process releasing two net molecules of ATP and two of NADH. The expression of glycolytic genes is regulated by HIF-1 α .

Hypoxia-inducible factor 1-alpha (HIF-1 α): a subunit of the heterodimeric transcription factor HIF-1 that, together with HIF-1 β , regulates the cellular response to hypoxia by inducing the transcriptional activation of genes involved in glycolysis.

phagocytophilum to infect tick hemocytes, which are required for the migration of *A. phagocytophilum* from the gut to the salivary glands [22]. Isolation and *in vitro* culture of tick hemocytes has remained elusive. Therefore, the study of metabolism in tick hemocytes in response to *A. phagocytophilum* infection has been based on the *I. scapularis* ISE6 cell line. Based on the transcriptional response to *A. phagocytophilum*, ISE6 cells were proposed to constitute a model of tick hemocytes involved in pathogen infection and immune response [23,24]. Comparison of the metabolic profile of M1 macrophages with that of *A. phagocytophilum*-infected ISE6 cells suggests that the metabolic profile of tick hemocytes in response to pathogen infection is very different to that of inflammatory M1 macrophages (Figure 1, Key Figure). Two major differences can be summarized: (i) ISE6 cells infected by *A. phagocytophilum* do not display the aerobic glycolysis observed in mammalian M1 macrophages, and (ii) **hypoxia-inducible factor 1-alpha (HIF-1 α)** activation and stabilization in ISE6 cells is not dependent on succinate accumulation and aerobic glycolysis (Figure 1). This suggests that the metabolic reprogramming associated with M1 macrophage activation does not occur in tick hemocytes infected with *A. phagocytophilum*. Interestingly, while 2-deoxyglucose (2DG), an inhibitor of **glycolysis**, inhibits HIF-1 α activation in M1 macrophages [25], this compound has no effect on HIF-1 α activation in ISE6 cells [26]. Instead of aerobic glycolysis and succinate accumulation, HIF-1 α activity in *A. phagocytophilum*-infected ISE6 cells appears to be mediated by activation of the phosphoinositide-3-kinase (PI3K) pathway (Figure 1). In agreement with this, the PI3K inhibitor LY294002 decreased HIF-1 α DNA binding activity in infected ISE6 cells [26]. A metabolic property shared by M1 macrophages and ISE6 cells is, however, an overall decrease in **(TCA cycle)** activity upon bacterial infection [25,26].

The absence of immunogenic molecules, such as LPS and diaminopimelic-type peptidoglycans (DAP-PGNs), in *A. phagocytophilum* and other bacteria transmitted by *I. scapularis* was previously suggested to contribute to vector tolerance to tick-borne pathogens [7]. The lack of these immune activators in tick-borne pathogens may help these microbes to colonize ticks without being eliminated [7]. Interestingly, lipids 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphoglycerol (POPG) and 1-palmitoyl-2-oleoyl diacylglycerol (PODAG), found in *A. phagocytophilum*, activate the immune deficiency (IMD) pathway in ticks [27]. Immune priming with POPG and PODAG reduces the *A. phagocytophilum* burden in infected ISE6 cells [27]. This shows that tolerance to infection can be associated with immunity and resistance to tick-borne pathogens in ticks. This is expected, considering that unchecked bacterial multiplication within host cells could affect vector, and therefore tick-borne pathogen, survival. Remarkably, recent research shows that IMD is involved in glucose and lipid metabolism regulation in *Drosophila* [28], and constitutive activation of IMD induces a significant reduction in the expression of genes involved in glycolysis, **gluconeogenesis**, and the TCA cycle in flies [29]. Whether IMD activation by *A. phagocytophilum* regulates tick hemocyte metabolism remains to be tested.

Molecular Mechanisms Involved in the Modulation of Tick Metabolism by Tick-borne Pathogens

Modulation of tick metabolism by tick-borne pathogens results mainly from the regulation of the levels of metabolic enzymes. Although it remains a challenge to explain how pathogen infection modifies the level of thousands of tick genes and hundreds of tick proteins [30], the molecular mechanisms behind this have been partially characterized using the model bacterium *A. phagocytophilum*. To induce transcriptional reprogramming in infected cells, *A. phagocytophilum* hijacks the transcriptional regulatory machinery of ticks by inducing the activation of signaling pathways [31] and the recruitment of transcription factors [32],

Oxaloacetate (OAA): a central metabolite produced in the TCA cycle that takes part in gluconeogenesis, amino acid metabolism, and fatty acid synthesis among other pathways. In addition, OAA and PEP result from anaplerotic reactions that keep the metabolic flux of glycolysis and gluconeogenesis.

Phosphoenolpyruvate (PEP): a key metabolite that stands at the crossroad between gluconeogenesis, glycolysis, and the TCA cycle and can be produced by the activity of two enzymes: enolase and PEPCK.

Phosphoenolpyruvate carboxykinase (PEPCK): PEPCK catalyzes the first committed step in gluconeogenesis, in which OAA is converted into PEP. There are two isoforms of the enzyme: mitochondrial (PEPCK-M) and cytosolic (PEPCK-C). PEPCK-C is a key regulatory enzyme of glyceroneogenesis and gluconeogenesis.

Tricarboxylic acid cycle (TCA cycle): a central route for oxidative phosphorylation in cells. In the TCA cycle, acetyl-CoA is oxidized to carbon dioxide, and reducing factors (NADH and FADH₂) – that contribute to the synthesis of ATP during oxidative phosphorylation – are produced. The TCA cycle is a crossroad of cellular metabolism in which carbohydrate, fatty acids, and amino acid metabolisms are interconnected.

Table 1. Tick Metabolic Pathways Affected by Tick-borne Pathogens

Metabolic pathways	Experimental system (<i>in vitro</i> vs. <i>in vivo</i>)	Affected by (name of pathogen species)	Effect on tick metabolism ^a	Refs
Glycolysis	ISE6 tick cells	<i>Anaplasma phagocytophilum</i>	Decreased levels and activity of HXK Increased levels of PEP Decreased levels of glucose and lactate	[26,43] [41] [23]
	<i>Ixodes scapularis</i> adults midguts		Upregulation of genes involved in glycolysis (except <i>hxx</i> and <i>pfk</i>) Upregulation of genes <i>hif-1α</i> and <i>hif-1β</i> Increased protein levels of glycolytic enzymes (except iPGM, PK, and PFK) and glucose transporters GLUT1A and SGLT2	[26]
	<i>Ixodes scapularis</i> adults salivary glands		Upregulation of genes involved in glycolysis (except <i>hxx</i> , <i>pfk</i> , <i>ald</i> , <i>pgk1</i> , and <i>eno1</i>) Increased protein levels of glycolytic enzymes (except HXK, PFK and ALDA) and glucose transporter GLUT1A	[26]
Gluconeogenesis	ISE6 tick cells	<i>Anaplasma phagocytophilum</i>	Decreased levels of PEPCK-C and glucose Increased protein levels of PEPCK-M	[23] [26]
	<i>Ixodes scapularis</i> adults midguts		Upregulation of genes involved in gluconeogenesis Decreased levels of gluconeogenesis enzymes (except FBP)	[26]
	<i>Ixodes scapularis</i> adults salivary glands		Decreased levels of gluconeogenesis enzymes	
Glyceroneogenesis	ISE6 tick cells	<i>Anaplasma phagocytophilum</i>	Decreased levels of glycerol-3-phosphate Increased levels of PEP, which may inhibit the enzyme TPI	[26]
	<i>Ixodes scapularis</i> adults midguts		Upregulation of genes <i>tpi</i> and <i>gpdh</i> Increased and decreased protein levels of TPI and GPDH, respectively	
	<i>Ixodes scapularis</i> adults salivary glands		Upregulation of gene <i>tpi</i> Increased protein levels of TPI and GPDH	
	<i>Ixodes scapularis</i> nymphs ^b	<i>Borrelia mayonii</i>	Decreased levels of glycerol-3-phosphate	[54]
Lipids	<i>Rhipicephalus microplus</i> adults fat body	<i>Metarhizium anisopliae</i>	Increased levels of cholesterol ester and triacylglycerol Decreased levels of free cholesterol	[51]
	<i>Ixodes scapularis</i> nymphs ^b	<i>Borrelia mayonii</i>	Decreased levels of cholesterol, ethanolamine phosphate, and ethanolamine	[54]
Amino acids	ISE6 tick cells	<i>Anaplasma phagocytophilum</i>	Decreased levels of alanine, glycine, glutamine, glutamate, and methionine Increased levels of serine, proline, tyrosine, and phenylalanine Tyrosine is used as fuel to synthesize PEP	[23,41]
			Upregulation of gene <i>oatp</i> Xanthurenic acid acts as a cofactor for unknown transcription factor(s) that control the expression of <i>oatp</i> KAT regulates the expression of <i>oatp</i>	[33]
	<i>Ixodes scapularis</i> adults midguts and salivary glands		Changes in the levels of genes and proteins involved in amino acid synthesis and degradation	[41]
	<i>Ixodes scapularis</i> nymphs salivary glands		Upregulation of genes <i>oatp</i> and <i>kat</i> Xanthurenic acid acts as a cofactor for unknown transcription factor(s) that control the expression of <i>oatp</i>	[33]

Table 1. (continued)

Metabolic pathways	Experimental system (<i>in vitro</i> vs. <i>in vivo</i>)	Affected by (name of pathogen species)	Effect on tick metabolism ^a	Refs
	<i>Ixodes scapularis</i> nymphs ^b	<i>Borrelia mayonii</i>	Decreased levels of alanine, isoleucine, leucine, lysine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, valine, glutamate, glutamine, and glycine	[54]
		<i>Borrelia burgdorferi</i>	Increased levels of tryptophan and glutamine Decreased levels of alanine, isoleucine, leucine, phenylalanine, proline, threonine, valine, and glycine	
Redox	BME26 tick cells	<i>Anaplasma marginale</i>	Downregulation of pro-oxidant genes and upregulation of antioxidant genes	[67]
	<i>Ixodes scapularis</i> adults midguts and salivary glands	<i>Anaplasma phagocytophilum</i>	Significant changes in the levels of genes and proteins of hemoglobinolytic enzymes affecting host hemoglobin levels in tick tissues	[18]

^aThe full names of enzymes are in the main text, except for phosphoglycerate kinase 1 (PGK1), enolase (ENOL), glycerol-3-phosphate dehydrogenase (GPDH) and triosephosphate isomerase (TPI).

^bOnly the levels of tick metabolites in response to *Borrelia* spp. infection at day 4 of feeding were considered [54].


metabolites [33], and histone-modifying enzymes [34] (Figure 2). *A. phagocytophilum* may also induce the degradation of tick proteins by the tick proteasome which, in turn, decreases the levels of targeted metabolic enzymes [23]. In addition, *A. phagocytophilum* infection induces a high proportion of alternatively spliced transcript events (ASEs), which increases the predicted gene ontology processes related to lipids, lipoproteins, and cholesterol/sterol metabolism and other processes in human HL-60 cells infected by this pathogen [35]. The occurrence of ASEs has not been tested in *A. phagocytophilum*-infected ticks or tick cell lines yet. However, there is no reason to rule out the hypothesis that ASEs also occur in ticks in response to *A. phagocytophilum* infection.

Activation of tick signaling pathways by *A. phagocytophilum* involves the G-protein-coupled-receptor (GPCR) that stimulates *Ixodes* p21-activated kinase (IPAK1) through the activation of PI3K, which results in actin-mediated signaling [31]. Activation of IPAK1 results in actin phosphorylation and accumulation of actin in the nucleus which mediates binding of RNA polymerase II (RNAPII) and TATA box-binding protein (TBP) to the promoter of the gene *salp16*, inducing its expression [31,36]. PI3K activation was also suggested to induce the expression of the genes *hif-1 α* , *hif-1 β* and acetyltransferase p300 upon *A. phagocytophilum* infection in tick cells [26] (Figure 2). Mobilization of HIF-1 α , HIF-1 β , and p300 at HIF target genes may induce HIF-mediated transcriptional activation of glycolytic genes [26].

Among the transcription factors (TFs) activated by *A. phagocytophilum* infection is the tick transcriptional activator protein-1 (AP-1) which induces the expression of the genes *iatgp* (*I. scapularis* antifreeze glycoprotein) and *kat* (kynurenine aminotransferase) in infected cells [32]. KAT is an enzyme involved in the production of the tryptophan metabolite xanthurenic acid [33]. Results suggests that xanthurenic acid acts as a cofactor of an uncharacterized TF that regulates the expression of one organic anion-transporting polypeptide (*oatp*) gene in *A. phagocytophilum*-infected ISE6 cells [33]. The upregulation of *oatp* and *kat* was proposed to be associated with increased levels of intracellular xanthurenic acid, which in turn favors *A. phagocytophilum* replication in tick salivary glands [33]. Conversely, RNAi-mediated silencing of *oatp* and *kat* expression decreases *A. phagocytophilum* levels in tick cells [33]. A general mechanism is then proposed by which *A. phagocytophilum* nucleomodulins recruit tick metabolites and TFs to activate target genes (Figure 2).


Key Figure

Metabolic Profiles of Tick Hemocytes and Mammalian Proinflammatory M1 Macrophages



Trends in Parasitology

Figure 1. The figure displays the metabolic profile of *Anaplasma phagocytophilum* (bacteria without LPS)-infected ISE6 cells, a model of tick hemocytes, compared to that of mammalian M1 macrophages in response to other intracellular bacteria with LPS. High levels of hexokinase (HXK, green) and phosphofruktokinase (PFK, green) are associated with high aerobic glycolysis and increased lactate production in mammalian M1 macrophages [25]. The levels and activity of tick HXK decreased (red), and the levels of tick PFK did not change (black) in response to infection which was associated with lower levels of lactate in infected ISE6 cells [23,26,43]. HIF-1 α activation and stabilization (green), that enhances the expression of glycolytic genes, is mediated by aerobic glycolysis and succinate accumulation from glutamine in M1 macrophages [25]. In consequence, blocking aerobic glycolysis with 2-deoxyglucose (2DG) inhibits HIF-1 α activation in these macrophages [25]. In contrast, 2DG does not inhibit HIF-1 α activation (green) in ISE6 cells, an effect that is achieved by LY294002, an inhibitor of the phosphoinositide-3-kinase (PI3K) pathway [26]. This suggests that HIF-1 α activity in tick hemocytes infected with *A. phagocytophilum* is regulated by PI3K and not aerobic glycolysis. M1 macrophages and ISE6 cells share a low TCA cycle activity [25,26]. Vertical arrows represent decrease (red) and increase (green) in metabolite levels. LPS, lipopolysaccharide.



Trends in Parasitology

Figure 2. Molecular Mechanisms Potentially Implicated in the Modulation of Tick Metabolism by the Model Pathogen *Anaplasma phagocytophilum*. Molecular mechanisms proposed to participate in the transcriptional reprogramming and proteome modulation induced by *A. phagocytophilum* infection in tick cells. Firstly, *A. phagocytophilum* activates the phosphoinositide-3-kinase (PI3K) signaling pathway in a G-protein-coupled-receptor (GPCR)-dependent manner to induce the expression of *salp16* [31], and hypoxia-inducible factor (HIF) *hif-1α* in ticks [26] (sequential steps and tick molecular factors colored in violet). Secondly, *A. phagocytophilum* recruits tick transcription factors (TFs), such as transcriptional activator protein-1 (AP-1) [32] and metabolites, such as xanthurenic acid [33], to regulate the expression of the metabolic genes *oatp* [33] and *kat* [32] (sequential steps and tick molecular factors colored in blue). Other tick metabolic genes may be regulated by the same mechanism in which we propose that *A. phagocytophilum* uncharacterized nucleomodulins may participate in the recruitment of tick TFs. Thirdly, the recruitment of histone deacetylase 1 (HDAC1) to the promoters of target genes, through the activity of nucleomodulins, may contribute to the repression of tick genes, for example, fructose-1,6-bisphosphatase (*fbp*) (sequential steps and tick molecular factors colored in green). Lastly, although the mechanism is unknown, *A. phagocytophilum* also targets tick proteins to the proteasome which, in turn, may result in the under-representation of targeted metabolic enzymes [23]. Abbreviations: IPAK1, *Ixodes* p21-activated kinase; RNAPII, RNA polymerase II; TBP, TATA box-binding protein; p300, acetyltransferase p300. Closed and open chromatin are represented as red and green double-stranded DNA, respectively.

The capacity of *A. phagocytophilum* to downregulate gene expression in mammalian neutrophils was associated with host histone deacetylase 1 (HDAC1) recruitment to the promoters of target genes by the bacterial nucleomodulin ankyrin repeat protein Anka [37]. Upon *A. phagocytophilum* infection, Anka is secreted through T4SS, enters the granulocyte nucleus, binds stretches of AT-rich DNA, and recruits host HDAC1 to its binding sites. *A. phagocytophilum* infection also induces HDAC1 binding and deacetylation of histone H3 (H3) which, in turn, results in downregulation of many host defense genes, including *CYBB*, which encodes a component (i.e., NOX2) of the NADPH oxidase [37]. Downregulation of *CYBB* reduces

superoxide anion production by NADPH oxidase which is essential for *A. phagocytophilum* survival in neutrophils [37]. Homologs of mammalian HDAC1 and H3 were identified in the *I. scapularis* genome [34]. Whether *A. phagocytophilum* AnkA is secreted and recruits HDAC1 to regulate gene transcription in infected ticks is currently unknown. However, *Ixodes* HDAC1 is over-represented in *A. phagocytophilum*-infected tick salivary glands, and chemical inhibition of this protein decreases the *A. phagocytophilum* burden in tick cells [34]. This shows that the expression and activity of HDAC1 is critical for *A. phagocytophilum* survival in both mammalian and tick cells. Notably, in mammalian cells, HDAC1 controls the expression of the metabolic enzyme fructose-1,6-bisphosphatase (FBP), a rate-limiting enzyme in gluconeogenesis [38]. Interestingly, high HDAC1 protein levels [34] were associated with low FBP protein levels in *A. phagocytophilum*-infected tick salivary glands [26]. Conversely, low HDAC1 protein levels [34] correlated with high mRNA and protein levels of *fbp* and FBP, respectively, in the *A. phagocytophilum*-infected tick midgut [26]. The negative correlation between the protein levels of HDAC1 and that of *fbp* and FBP strongly suggests that HDAC1 may be recruited by *A. phagocytophilum* nucleomodulins to act as a repressor of *fbp* expression in tick salivary glands (Figure 2). A decrease in FBP protein levels may contribute to the inhibition of gluconeogenesis that *A. phagocytophilum* induces in infected tick cells (see below) [23].

Tissue-specific Changes in Tick Carbohydrate Metabolism in Response to Infection

Glucose is the most abundant carbohydrate in the hemolymph of adult ticks and it is required in tick embryonic development [39]. The regulation of glucose metabolic pathways in ticks is essential for the survival of *A. phagocytophilum* in the vector [23,26]. Notably, *A. phagocytophilum* infection inhibits gluconeogenesis by decreasing the protein levels of **phosphoenolpyruvate carboxy kinase-cytosolic** (PEPCK-C) which results in lower levels of glucose in *A. phagocytophilum*-infected ISE6 cells [23,26]. PEPCK-C is also a key regulatory enzyme in **glyceroneogenesis** [40] and thus, low levels of PEPCK-C were associated with reduced glycerol-3-phosphate production in infected tick cells [41]. Low PEPCK-C activity and decreased glucose anabolism inhibits apoptosis, which is essential for *A. phagocytophilum* survival in tick cells and tissues [23,30]. Interestingly, glycerol-3-phosphate levels were also reduced in tick nymphs infected by *Borrelia mayonii* (Table 1). Low levels of glycerol-3-phosphate may have a significant impact in fatty acids metabolism, a poorly explored facet in tick-pathogen interactions.

Tissue-specific differential regulation of apoptosis pathways was observed in adult female midgut and salivary glands in response to *A. phagocytophilum* infection [42]. Likewise, the transcriptional reprogramming induced by *A. phagocytophilum* affects the levels of carbohydrate metabolism genes and proteins in a tissue-specific manner [26]. For example, the protein levels of all glycolytic enzymes increase in *A. phagocytophilum*-infected tick midguts, except for the cofactor-independent phosphoglycerate mutase (iPGM) and pyruvate kinase (PK); despite gene upregulation, these two proteins were under-represented in this tissue [26]. In addition, gene and protein levels of phosphofructokinase (PFK) did not change in tick midguts in response to infection [26]. However, in salivary glands, the glycolysis enzymes fructose-bisphosphate aldolase A (ALDA), PFK, and hexokinase (HXK) were under-represented while the rest of the glycolytic enzymes were over-represented. To accurately assess the metabolic impact of the differential protein representation in tick midguts and salivary glands, a tissue-specific metabolomics approach is required. However, the lower levels of PK and HXK, key regulatory enzymes in glycolysis, in infected tick midguts and salivary glands, respectively, suggest that the production of glycolytic intermediates may be affected in a tissue-specific manner. It would also be important to assess the enzymatic activity of key enzymes which may vary in *A. phagocytophilum*-infected cells and tissues. For example, it was reported that the

activity of HXK decreases more than twofold in *A. phagocytophilum*-infected ISE6 cells [43]. The transcriptional activation of tick glycolytic genes was proposed to be mediated by tick HIF-1 α [26], which contains structural domains conserved in model organisms, and the gene *hif-1 α* was significantly upregulated in infected tick midguts [26]. Another important gene targeted by HIF-1 α is pyruvate dehydrogenase kinase 1 (*pdk1*) which encodes the protein PDK1 that controls the enzymatic activity of pyruvate dehydrogenase E1 (PDE1) which transforms pyruvate into **acetyl-CoA**, the first step in the TCA cycle. PDK1 mRNA and protein levels were higher in *A. phagocytophilum*-infected tick salivary glands, suggesting that this pathogen inhibits the TCA cycle in infected salivary glands [26].

Crosstalk between Tick Amino Acid and Carbohydrate Metabolism during Pathogen Infection

Several recent studies have shown that amino acids are central to host–pathogen metabolic interaction [3,23,41,44]. When digested, the protein-rich diet of ticks generates large amounts of amino acids. Therefore, amino acid metabolism is of great interest in ticks. For example, tyrosine-associated toxicity is lethal for blood-sucking arthropods, which points to the significance of tyrosine degradation pathways. Inhibition of tyrosine aminotransferase (TAT) and 4-hydroxyphenylpyruvate dioxygenase (HPPD), the first two enzymes of the phenylalanine/tyrosine degradation pathway, caused the death of mosquitoes, kissing bugs, and ticks after a blood meal [45].

Infection by *A. phagocytophilum* rewires the network of tick vector cell processes and changes the relative importance of some biological pathways, including those involved in metabolism [46]. This is to be expected, considering that *A. phagocytophilum* has a small genome and cannot actively carry out several metabolic processes, and therefore requires and hijacks tick cell resources for survival within the vector [47]. For example, out of 20 amino acids, the *A. phagocytophilum* genome encodes only those enzymes responsible for the biosynthesis of four amino acids (proline, glutamine, glycine, and aspartate), and the glycolysis enzymes are reduced to those that produce glyceraldehyde-3-phosphate and dihydroxyacetone phosphate (DHAP) from **phosphoenolpyruvate (PEP)** [47]. As a consequence, *A. phagocytophilum* changes the levels of amino acid metabolism enzymes [41], which, in turn, affects the intracellular levels of several amino acids and metabolic intermediates [23,33]. Interestingly, tick cells use tyrosine as a fuel to synthesize PEP during early *A. phagocytophilum* infection [41]. This process is achieved by activating the tyrosine/**oxaloacetate** (OAA)/PEPCK-M/PEP node by which *A. phagocytophilum* infection may decrease the tyrosine pool, which, in turn, may protect ticks against tyrosine-induced toxicity *in vivo* [41].

Tick-borne flaviviruses are known to affect protein processing and amino acid metabolism in tick cells [48,49]. Particularly, infection by Langat virus (LGTV) increased protein levels of acetyl-CoA acetyltransferase 1 (ACAT1), aldehyde dehydrogenase (DP5CD), glutamate dehydrogenase (GLUD1), and fumarylacetoacetase (FAH) in ISE6 cells [49]. Knockdown of the gene *acat1* was associated with decreased LGTV genome replication and LGTV particle release, while knockdown of the gene *fah* decreased only LGTV particle release in ISE6 cells [48]. Interestingly, FAH catalyzes the last step in tyrosine degradation, suggesting that flavivirus infection may, similarly to *A. phagocytophilum*, contribute to tyrosine detoxification in ticks.

Tick Lipid Metabolism Is Exploited by Different Tick-associated Microorganisms

Starvation and desiccating conditions that ticks have to endure during off-host periods affect the energy reserves of the tick [50]. For example, starvation of *Dermacentor variabilis* ticks in

optimal humidity conditions is known to result in up to 60% losses in lipid reserves [50]. The parietal fat body, the main lipid reserve organ for triacylglycerol and cholesterol, is gaining increasing interest for the study of infections in insects and ticks [51,52]. However, lipid metabolism signaling pathways are still rarely studied in ticks.

Fatty acids play important roles as energy sources for several microorganisms, including pathogenic bacteria [53]. If a pathogen can prevent the use of energy reserves by the host, this will affect its immune system, thereby promoting pathogen infection and persistence. *B. burgdorferi* is not able to synthesize fatty acids and cholesterol, thus relying on the vertebrate host cell or the blood meal of the tick for lipid supply [54]. Similarly, host cholesterol and other lipids are required by *A. phagocytophilum* for infection and multiplication in human cells [55]. Perilipin (PLIN), a major adipocyte lipid droplet-associated phosphoprotein that plays a central role during lipolysis and cholesterol synthesis, is differentially expressed in HL-60 cells in response to infection with *A. phagocytophilum* [56]. The bacteria are able to alter PLIN levels in infected cells, thus modifying the lipid metabolism of the host to facilitate bacterial infection [56]. This phosphoprotein is likely to play a similar role in the metabolism of the tick, although its role has not been characterized so far.

Other obligate intracellular pathogens, such as viruses, need to hijack the cellular machinery for the synthesis of lipids to complete their replication. The flavivirus life cycle is very closely associated with host cell lipids, and these viruses can rearrange intracellular membranes and reorganize the host cell lipid metabolism of infected cells to ensure viral replication [57]. Dengue virus infection can alter metabolic pathways in humans, such as fatty acid biosynthesis and β -oxidation [58]. Exosomes are small extracellular vesicles that appear to play a central role in cell-to-cell communication [58]. Exosomes transport bioactive molecules, such as messenger RNAs (mRNAs) and microRNAs (miRNAs) that can be transferred in active form to adjacent cells or to distant organs, thus being important vehicles for metabolic cross-talk [59]. Recent research has shown that flaviviruses use arthropod-derived exosomes secreted by tick cells as a means for viral RNA and metabolite transmission from the vector, and the vertebrate exosomes for dissemination within the host [60]. The knockdown of transcripts for a glycosylphosphatidylinositol (GPI)-anchored protein (VNN) was associated with decreased LGTV genome replication and LGTV release in ISE6 tick cells [48]. VNN is involved in coenzyme-A (CoA) metabolism [61], which is critical for lipid metabolism.

Fungal infections can also modulate the lipid metabolism of the host [62]. Recent research with different isolates of *Metarhizium anisopliae* showed that infection of *Rhipicephalus microplus* ticks with entomopathogenic fungi can modulate the lipid content, thus inducing changes in important metabolic pathways [62]. However, lipid metabolism modulation during *M. anisopliae* infection in ticks was independent of AMP-activated protein kinase (AMPK) and extracellular signal-regulated protein kinase (ERK) pathways [62] which are the major pathways involved in the breakdown of triacylglycerol into fatty acids in mammals [63,64]. This suggested that modulation of tick lipid metabolism by *M. anisopliae* occurs through an alternative and uncharacterized lipolysis pathway.

Concluding Remarks

Interactions between ticks, hosts, and pathogens involve many metabolic pathways, such as carbohydrate, protein, lipid and redox pathways, that are affected with the production of either conflict or cooperation between them. These interactions at the metabolic level lead to mutual beneficial effects in tick–host–pathogen molecular interactions. Among them, amino

Outstanding Questions

Do ticks use iron metabolism and nutritional immunity to control microbial infection? What mechanisms use tick-borne pathogens to overcome nutritional immunity in ticks?

Does modulation of tyrosine metabolism by *Anaplasma phagocytophilum* increase tick survival to tyrosine-induced toxicity?

What is the contribution of tick epigenetics to tick metabolism regulation? Is histone deacetylation a central mechanism for metabolic gene repression in ticks? Do *A. phagocytophilum* nucleomodulins recruit tick histone deacetylases to repress tick metabolic genes?

What molecular mechanisms account for lipid metabolism modulation by *A. phagocytophilum* and *Borrelia burgdorferi* in ticks? Why does *A. phagocytophilum* and *B. burgdorferi* reduce, significantly, tick glycerol-3-phosphate levels? What is the role of this metabolite in tick–pathogen metabolic interactions?

acids are central to the host–pathogen metabolic interaction. Nevertheless, synthesis and degradation of amino acids is narrowly connected to carbohydrate metabolism, and the later to lipid metabolism.

Despite recent advances in the study of tick–host–pathogen molecular interactions, most of the pathways and molecules involved in these interactions are not fully characterized. Furthermore, the regulatory mechanisms affecting the metabolic pathways affected by these interactions are only partially characterized (see Outstanding Questions). Future directions should include the application of omics technologies together with data integration algorithms [46] to the investigation of tick interactome, regulome, miRNAome, epigenome, and metabolome and their interactions to advance the metabolic perspective on tick–pathogen interactions.

Identification of the metabolic adaptations during tick–pathogen interactions that support tick survival, spread, and pathogen infection and transmission could provide new therapeutic targets for the prevention and control of tick-borne diseases. Studies of the interactions between the pathogen and the host lipid metabolism can help to identify potential targets for the development of vaccines or treatments against health-threatening pathogens. These strategies could include the inhibition of lipid biosynthesis. However, manipulation of this major metabolic pathway can also be detrimental for the host. For instance, treatment with Lovastatin, an inhibitor of cholesterol synthesis, appears to delay infection and increases survival rates in AG129 mice infected with dengue virus serotype 2 while it also tends to increase viremia, which could affect disease progression [65]. A better understanding of the interactions of pathogens with the host cell lipid metabolism and other metabolic pathways could provide new therapeutic targets to control infections by tick-borne pathogens. Furthermore, as recently proposed, the application of machine learning and big data analysis to the tick–host–pathogen interaction datasets may also lead to the high-throughput identification of candidate vaccine protective antigens [66].

Acknowledgments

PA was funded by the Junta de Comunidades of Castilla-La Mancha (Spain).

References

- Ganeshan, K. and Chawla, A. (2014) Metabolic regulation of immune responses. *Annu. Rev. Immunol.* 32, 609–634
- Green, D.R. *et al.* (2014) Cell biology: Metabolic control of cell death. *Science*, 345, 1250256. 10 February 2016
- Eisenreich, W. *et al.* (2015) Metabolic adaptations of intracellular bacterial pathogens and their mammalian host cells during infection ('pathometabolism'). *Microbiol. Spectr.* 3, 27–58
- de la Fuente, J. *et al.* (2016) Tick–host–pathogen interactions: conflict and cooperation. *PLoS Pathog.* 12, e1005488
- Cabezas-Cruz, A. *et al.* (2017) Tick–pathogen ensembles: do molecular interactions lead ecological innovation? *Front. Cell. Infect. Microbiol.* Published online March 13, 2017. <https://doi.org/10.3389/fcimb.2017.00074>
- Busby, A.T. *et al.* (2012) Expression of heat shock proteins and subolesin affects stress responses, *Anaplasma phagocytophilum* infection and questing behaviour in the tick, *Ixodes scapularis*. *Med. Vet. Entomol.* 26, 92–102
- Shaw, D.K. *et al.* (2018) Vector immunity and evolutionary ecology: the harmonious dissonance. *Trends Immunol.* 39, 862–873
- de la Fuente, J. *et al.* (2015) Flying ticks: anciently evolved associations that constitute a risk of infectious disease spread. *Parasit. Vectors.* 8, 538
- de la Fuente, J. *et al.* (2016) *Anaplasma phagocytophilum* uses common strategies for infection of ticks and vertebrate hosts. *Trends Microbiol.* 24, 173–180
- Guila-Nuss, M. *et al.* (2016) Genomic insights into the *Ixodes scapularis* tick vector of Lyme disease. *Nat. Commun.* 7, 10507
- de la Fuente, J. *et al.* (2017) Tick–pathogen interactions and vector competence: identification of molecular drivers for tick-borne diseases. *Front. Cell. Infect. Microbiol.* Published online April 7, 2017. <https://doi.org/10.3389/fcimb.2017.00114>
- Núñez, G. *et al.* (2018) Innate nutritional immunity. *J. Immunol.* 201, 11–18
- Galay, R.L. *et al.* (2015) Iron metabolism in hard ticks (Acari: Ixodidae): the antidote to their toxic diet. *Parasitol. Int.* 64, 182–189
- Moumène, A. *et al.* (2018) Iron starvation conditions upregulate *Ehrlichia ruminantium* type IV secretion system, *tr1* transcription factor and *map1* genes family through the master regulatory protein ErxR. *Front. Cell. Infect. Microbiol.* Published online January, 2018. <https://doi.org/10.3389/fcimb.2017.00535>
- Kahlon, A. *et al.* (2013) *Anaplasma phagocytophilum* Asp14 is an invasin that interacts with mammalian host cells via its C terminus to facilitate infection. *Infect. Immun.* 81, 65–79
- Villar, M. *et al.* (2015) Identification and characterization of *Anaplasma phagocytophilum* proteins involved in infection of the tick vector, *Ixodes scapularis*. *PLoS One*, Published online <https://doi.org/10.1371/journal.pone.0137237>
- Troxell, B. *et al.* (2012) *Borrelia burgdorferi*, a pathogen that lacks iron, encodes manganese-dependent superoxide dismutase

- essential for resistance to streptonigrin. *J. Biol. Chem.* 287, 19284–19293
18. Villar, M. *et al.* (2016) The intracellular bacterium *Anaplasma phagocytophilum* selectively manipulates the levels of vertebrate host proteins in the tick vector *Ixodes scapularis*. *Parasit. Vectors*, 9, 467
 19. Sachla, A.J. *et al.* (2014) The crimson conundrum: heme toxicity and tolerance in GAS. *Front. Cell. Infect. Microbiol.* Published online November 5, 2014. <https://doi.org/10.3389/fcimb.2014.00159>
 20. Price, J.V. and Vance, R.E. (2014) The macrophage paradox. *Immunity*, 41, 685–693
 21. Hajdušek, O. *et al.* (2013) Interaction of the tick immune system with transmitted pathogens. *Front. Cell. Infect. Microbiol.* Published online July 16, 2013. <https://doi.org/10.3389/fcimb.2013.00026>
 22. Liu, L. *et al.* (2011) *Ixodes scapularis* salivary gland protein P11 facilitates migration of *Anaplasma phagocytophilum* from the tick gut to salivary glands. *EMBO Rep.* 12, 1196–1203
 23. Villar, M. *et al.* (2015) Integrated metabolomics, transcriptomics and proteomics identifies metabolic pathways affected by *Anaplasma phagocytophilum* infection in tick cells. *Mol. Cell. Proteom.* 14, 3154–3172
 24. Alberdi, P. *et al.* (2016) Tissue-specific signatures in the transcriptional response to *Anaplasma phagocytophilum* infection of *Ixodes scapularis* and *Ixodes ricinus* tick cell lines. *Front. Cell. Infect. Microbiol.* Published online February 10, 2016. <https://doi.org/10.3389/fcimb.2016.00020>
 25. Nonnenmacher, Y. and Hiller, K. (2018) Biochemistry of proinflammatory macrophage activation. *Cell. Mol. Life. Sci.* 75, 2093–2109
 26. Cabezas-Cruz, A. *et al.* (2017) *Anaplasma phagocytophilum* infection subverts carbohydrate metabolic pathways in the tick vector, *Ixodes scapularis*. *Front. Cell. Infect. Microbiol.* Published online February 7, 2017. <https://doi.org/10.3389/fcimb.2017.00023>
 27. Shaw, D.K. *et al.* (2017) Infection-derived lipids elicit an immune deficiency circuit in arthropods. *Nat. Commun.* 8, 14401
 28. Kamareddine, L. *et al.* (2018) The *Drosophila* immune deficiency pathway modulates enteroendocrine function and host metabolism. *Cell. Metab.* 28, 449–462
 29. Davoodi, S. *et al.* (2018) The immune deficiency pathway regulates metabolic homeostasis in *Drosophila*. *bioRxiv*, Published online December 14, 2018. <https://doi.org/10.1101/253542>
 30. Aylón, N. *et al.* (2015) Systems biology of tissue-specific response to *Anaplasma phagocytophilum* reveals differentiated apoptosis in the tick vector *Ixodes scapularis*. *PLoS Genet.* 11, e1005120
 31. Sultana, H. *et al.* (2010) *Anaplasma phagocytophilum* induces actin phosphorylation to selectively regulate gene transcription in *Ixodes scapularis* ticks. *J. Exp. Med.* 207, 1727–1743
 32. Khanal, S. *et al.* (2018) Arthropod transcriptional activator protein-1 (AP-1) aids tick-rickettsial pathogen survival in the cold. *Sci. Rep.* 8, 11409
 33. Taank, V. *et al.* (2017) Human rickettsial pathogen modulates arthropod organic anion transporting polypeptide and tryptophan pathway for its survival in ticks. *Sci. Rep.* 7, 13256
 34. Cabezas-Cruz, A. *et al.* (2016) *Anaplasma phagocytophilum* increases the levels of histone modifying enzymes to inhibit cell apoptosis and facilitate pathogen infection in the tick vector *Ixodes scapularis*. *Epigenetics*, 11, 303–319
 35. Dumler, J.S. *et al.* (2018) Alternative splicing of differentiated myeloid cell transcripts after infection by *Anaplasma phagocytophilum* impacts a selective group of cellular programs. *Front. Cell. Infect. Microbiol.* Published online February 2, 2018. <https://doi.org/10.3389/fcimb.2018.00014>
 36. Sukumaran, B. *et al.* (2006) An *Ixodes scapularis* protein required for survival of *Anaplasma phagocytophilum* in tick salivary glands. *J. Exp. Med.* 203, 1507–1517
 37. Rennoll-bankert, K.E. *et al.* (2015) Chromatin-bound bacterial effector ankyrin A recruits histone deacetylase 1 and modifies host gene expression. *Cell. Microbiol.* 17, 1640–1652
 38. Yang, J. *et al.* (2017) Inhibiting histone deacetylases suppresses glucose metabolism and hepatocellular carcinoma growth by restoring FBP1 expression. *Sci. Rep.* 7, 43864
 39. Martins, R. *et al.* (2018) The dynamics of energy metabolism in the tick embryo. *Rev. Bras. Parasitol. Vet.* 27, 259–266
 40. Saponaro, C. *et al.* (2015) The subtle balance between lipolysis and lipogenesis: a critical point in metabolic homeostasis. *Nutrients*, 7, 9453–9474
 41. Cabezas-Cruz, A. *et al.* (2017) *Ixodes scapularis* tick cells control *Anaplasma phagocytophilum* infection by increasing the synthesis of phosphoenolpyruvate from tyrosine. *Front. Cell. Infect. Microbiol.* Published online August 17, 2017. <https://doi.org/10.3389/fcimb.2017.00375>
 42. Alberdi, P. *et al.* (2016) *Anaplasma phagocytophilum* manipulates host cell apoptosis by different mechanisms to establish infection. *Vet. Sci.* Published online July, 2016. <https://doi.org/10.3390/vetsci3030015>
 43. Aylón, N. *et al.* (2013) *Anaplasma phagocytophilum* inhibits apoptosis and promotes cytoskeleton rearrangement for infection of tick cells. *Infect. Immunity*. 81, 2415–2425
 44. Olive, A.J. and Sasseti, C.M. (2016) Metabolic crosstalk between host and pathogen: sensing, adapting and competing. *Nat. Rev. Microbiol.* 14, 221–234
 45. Sterkel, M. *et al.* (2016) Tyrosine detoxification is an essential trait in the life history of blood-feeding arthropods. *Curr. Biol.* 26, 2188–2193
 46. Estrada-Peña, A. *et al.* (2018) Use of graph theory to characterize human and arthropod vector cell protein response to infection with *Anaplasma phagocytophilum*. *Front. Cell. Infect. Microbiol.* Published online August 3, 2018. <https://doi.org/10.3389/fcimb.2018.00265>
 47. Dunning Hotopp, J.C. *et al.* (2006) Comparative genomics of emerging human ehrlichiosis agents. *PLoS Genet.* 2, e21
 48. Grabowski, J.M. *et al.* (2017) RNAi reveals proteins for metabolism and protein processing associated with Langat virus infection in *Ixodes scapularis* (black-legged tick) ISE6 cells. *Parasit. Vectors*, Published online January 13, 2017. <https://doi.org/10.1186/s13071-016-1944-0>
 49. Grabowski, J.M. *et al.* (2016) Changes in the proteome of Langat-infected *Ixodes scapularis* ISE6 cells: metabolic pathways associated with flavivirus infection. *PLoS Negl. Trop. Dis.* Published online February 9, 2016. <https://doi.org/10.1371/journal.pntd.0004180>
 50. Rosendale, A.J. *et al.* (2017) Dehydration and starvation yield energetic consequences that affect survival of the American dog tick. *J. Insect. Physiol.* 101, 39–46
 51. Angelo, I.C. *et al.* (2013) Neutral lipid composition changes in the fat bodies of engorged females *Rhipicephalus microplus* ticks in response to fungal infections. *Parasitol. Res.* 112, 501–509
 52. Charroux, B. and Royet, J. (2010) *Drosophila* immune response: From systemic antimicrobial peptide production in fat body cells to local defense in the intestinal tract. *Fly (Austin)*, 4, 40–47
 53. Barisch, C. and Soldati, T. (2017) Breaking fat! How mycobacteria and other intracellular pathogens manipulate host lipid droplets. *Biochimie*, 141, 54–61
 54. Hoxmeier, J.C. *et al.* (2017) Metabolomics of the tick–Borrelia interaction during the nymphal tick blood meal. *Sci. Rep.* 7, 44394
 55. Truchan, H.K. *et al.* (2013) Breaking in and grabbing a meal: *Anaplasma phagocytophilum* cellular invasion, nutrient acquisition, and promising tools for their study. *Microbes Infect.* 15, 1017–1025
 56. Manzano-Roman, R. *et al.* (2008) Expression of perilipin in human promyelocytic cells in response to *Anaplasma phagocytophilum* infection results in modified lipid metabolism. *J. Med. Microbiol.* 57, 159–163

57. Martín-Acebes, M.A. *et al.* (2016) Lipids and flaviviruses, present and future perspectives for the control of dengue, Zika, and West Nile viruses. *Prog. Lipid. Res.* 64, 123–137
58. Chotiwan, N. *et al.* (2018) Dynamic remodeling of lipids coincides with dengue virus replication in the midgut of *Aedes aegypti* mosquitoes. *PLoS Pathog.* 14, e1006853
59. Yu, X. *et al.* (2016) Exosomes as miRNA carriers: formation-function-future. *Int. J. Mol. Sci.* 17, 2028
60. Zhou, W. *et al.* (2018) Exosomes serve as novel modes of tick-borne flavivirus transmission from arthropod to human cells and facilitates dissemination of viral RNA and proteins to the vertebrate neuronal cells. *PLoS Pathog.* 14, e1006764
61. Kinoshita, T. and Fujita, M. (2016) Biosynthesis of GPI-anchored proteins: special emphasis on GPI lipid remodeling. *J. Lipid. Res.* 57, 6–24
62. Sá, F.A. *et al.* (2018) *Metarhizium anisopliae* s.l. modulation of lipid metabolism during tick infection is independent of AMPK and ERK pathways. *Parasitol. Res.* 117, 793–799
63. Jiao, P. *et al.* (2013) Hepatic ERK activity plays a role in energy metabolism. *Mol. Cell. Endocrinol.* 375, 157–166
64. Hardie, D.G. (2014) AMPK-sensing energy while talking to other signaling pathways. *Cell. Metab.* 20, 939–952
65. Martínez-Gutiérrez, M. *et al.* (2014) Lovastatin delays infection and increases survival rates in AG129 mice infected with dengue virus serotype 2. *PLoS One*, Published online February 21, 2014. <https://doi.org/10.1371/journal.pone.0087412>
66. de la Fuente, J. *et al.* (2018) High throughput discovery and characterization of tick and pathogen vaccine protective antigens using vaccinomics with intelligent Big Data analytic techniques. *Expert. Rev. Vaccines*, 17, 569–576
67. Kaili, S.P. *et al.* (2017) Immune-related redox metabolism of embryonic cells of the tick *Rhipicephalus microplus* (BM) in response to infection with *Anaplasma marginale*. *Parasit. Vectors*, E26 Published online December 19, 2017. <https://doi.org/10.1186/s13071-017-2575-9>